

Using gas geochemistry to delineate structural compartments and assess petroleum reservoir-filling directions: A Venezuelan case study

G. MÁRQUEZ^{a,*}, M. ESCOBAR^b, E. LORENZO^a, J.R. GALLEGÓ^c, R. TOCCO^d

^a Departamento de Ingeniería Minera, Mecánica y Energética, Universidad de Huelva, 21819 Huelva, Spain

^b CARBOZULIA S.A., Av. 2 No. 55-185, Casa Mene Grande & Universidad del Zulia, Maracaibo 4002 A, Venezuela

^c Departamento de Exploración y Prospección de Minas, Universidad de Oviedo, 33600 Mieres, Spain

^d REPSOL Exploración S.A., Paseo de la Castellana 280, 28046 Madrid, Spain

Abstract: Here we examined the light hydrocarbon and nitrogen content and isotopic signatures of eleven gaseous samples in order to evaluate lateral intra-reservoir continuity in a Venezuelan reservoir in the central area of Lake Maracaibo Basin. At least three single compartments, located in the northern-central and southern parts of the reservoir, are revealed by nitrogen concentrations showing clear step-like compositional breaks. The occurrence of step-breaks was also supported by the isotopic signature of individual hydrocarbon compounds in the range of C₁ to C₄ alkanes. Samples presented only slight differences in N₂ and hydrocarbon gas compositions within the central and northern parts of the reservoir, and therefore it was not possible to infer structural barriers in coherence with the geological section. Some oil bulk parameters corroborate gradual changes that provide additional information on the reservoir-filling history, thus suggesting that the lateral physical-chemical equilibrium of fluids was not reached in this reservoir.

Key-words: isotope geochemistry, reservoir compartments, hydrocarbon gas composition, nitrogen, C₁-C₄ alkanes.

1. Introduction

Organic geochemistry has played a key role in hydrocarbon exploration in recent decades. In particular, the application of gas geochemistry to reservoir studies and production engineering has attracted considerable attention (Kaufman et al., 1990; Bazan, 1998; Levaché et al., 2000; among others). Several case studies have reported the use of oil and gas compositional variations to investigate reservoir compartmentalization (i.e., Schoell and Jenden, 1993; Hwang et al., 1994; Larter and Aplin, 1994; Holba et al., 1996; Beeunas et al., 1999; Tocco and Alberdi, 1999).

Reservoir geology and engineering are being integrated by means of new analytical techniques that provide information on the natural complexity of oils. For example, a novel complementary method based on Fourier Transform Infra Red (FTIR) spectroscopy was recently proposed (Permanyer et al., 2002) to determine crude oil reservoir compartmentalization. In addition, time-lapse geochemistry (TLG) technology was used for two reservoirs in the deep-water Gulf of Mexico to locate and study petroleum barriers, thus yielding valuable information about reservoir behavior (Milkov et al., 2007). The study of low molecular-weight fractions ($\leq C_{15}$), and more recently gas fingerprints to evaluate the lateral and vertical continuity of reservoirs is based on the premise that light fractions from a single hydrocarbon column (continuous reservoir) have similar chemical compositions, whereas production from separate hydrocarbon columns (different sand/fault blocks) implies significantly different compositions (England et al., 1995). In this context, the use of gas isotopes to assess reservoir fluid heterogeneity has proved useful to further our understanding of reservoir connectivity (Rein and Shulz, 2007). The reasons for reservoir chemical and isotopic differences in discontinuous hydrocarbon columns are not fully understood and therefore a detailed study of each reservoir is required (Wavrek et al., 2001). Compositional similarities are thought to be the result of homogenization within the compartments after diffusion, convection (density-driven), and fluid-rock interactions have reached equilibrium (Smalley et al., 1994). England (2007) reviewed the applications of reservoir geochemistry from a reservoir engineering point of view and concluded that it is useful for addressing issues related to production allocation, reservoir compartmentalization, and the prediction of gravitational gradients. Beeunas et al. (1996) described in two reservoir sands without communication in the Gulf of Mexico a similar situation when compared to the present case study, determining carbon isotopic variations in methane, ethane, and propane.

It is also pertinent to note that nitrogen is one of the main non-hydrocarbon gas in many petroliferous basins (Jenden et al., 1988). Although nitrogen in natural gases is probably derived from organic matter

in the sedimentary column (Rohrback et al., 1983), there are other subsurface nitrogen sources such as dissolved air in groundwater, mantle-derived N₂ in areas of recent magmatic activity, and N₂ released from metasedimentary rocks during metamorphism (Gold and Held, 1987; Krooss et al., 1993). While N₂ concentrations have been used mostly to allocate gas mixtures in storage containers (Schoell and Jenden, 1993), molecular nitrogen concentration in gases can also be applied to reservoir characterization (Ballentine and Sherwood Lollar, 2002). Consequently, there is growing interest in modeling N₂ generation (Littke et al., 1993). To date, most nitrogen studies have been done in gas-source rocks, such as coal or source rock containing humic organic matter (Clayton, 1997; among others); however, nitrogen generation from marine organic matter has received less attention (i.e., Kotarba and Nagao, 2008).

Given the preceding considerations, here we integrated diverse geochemical data with the aim to elucidate lateral intra-reservoir continuity in a Venezuelan reservoir (central sector of Lake Maracaibo Basin). For this purpose, we applied a relatively recent approach to integrate information about the compositional signatures of the C₁-C₄ alkanes and molecular nitrogen with isotopic analyses of individual hydrocarbons in gaseous samples. Furthermore, API gravities, V/Ni ratios and sulfur contents in sampled oils are also provided to complement and contrast data obtained from gases.

2. Petroleum exploitation context

The Maracaibo Basin is located in north-western Venezuela (Figure 1a), an area comprising mainly the state of Zulia and also those of Falcón, Mérida, Táchira, and Trujillo. The Lake Maracaibo Basin

(containing 700 oil-producing fields after drilling about 16,500 wells) covers an area of approximately 50,000 km², with an estimated sediment volume of 250,000 km³ over a pre-Cretaceous basement (González de Juana et al., 1980).




Fig. 1. a) Map of Lake Maracaibo in northwestern part of Venezuela; b) and c), respectively, stratigraphic column and sample locations in the study area

The main petroleum source rocks in the study area are the Cenomanian-Campanian La Luna Formation (Erlich et al., 2000) and, to a lesser extent, the Aptian-Albian Apón Formation (Machiques Member), although other rock units also generate hydrocarbons (Talukdar et al., 1986). The Lake Maracaibo Basin occupies the entire lake area (12,870 km² with water to an average depth of 100 ft) and the

surrounding rolling hills. The major hydrocarbon accumulations are found in Eocene and Miocene deltaic sandstones (Talukdar and Marcano, 1994).

There are series of very large oilfields in the central sector of Lake Maracaibo Basin, including Lago-Lamar, Block I, Block II, Block V, and Ambrosio, among others. The stratigraphy of this sector consists of clastic sedimentary rocks of Miocene, Oligocene, Eocene, Paleocene and Cretaceous Age (see [Figure 1b](#)). The main reservoir rocks are the Eocene Misoa Formation, as well as the Miocene Lagunillas and La Rosa formations (Parnaud et al., 1995).

3. Methodology

Eleven natural gas samples from nine oil wells (labeled between W-01 and W-09) and two gas-condensate wells (W-10 and W-11) were collected in sealed glass bottles by means of metal separators connected to two sealed cylinders in parallel in the well-head. Additionally, ten oil samples (W-01 to W-09, and W-14) were also collected and stored in glass jars prior to analysis. In both cases, a proper seal prevented the loss of light compounds and other possible sample alterations. Figure 1c shows the sample collection sites in the study area. All these producing wells are within the boundary of the study reservoir's area located in the Lamasystem fault zone. A secondary normal faulting (represented by "I", see Figure 2) in this fault system would act as a sealing structure. This structural barrier must be considered, even though the displacement of the reservoir sands in the geological section does not seem to be high enough to interbed these sands with the over- and underlying beds in the stratigraphic framework. In the aforementioned section, each secondary fault is given a Roman numeral.




Fig. 2. North-south simplified cross-section of the study area. It includes from bottom to top the Misoa, Icotea, La Rosa, Lagunillas, Isnotu and Onia Fms.

Analyses of compounds in gaseous samples were carried out in an HP-5890 Series II gas chromatograph with electronic pneumatic control. The multidimensional gas chromatography system includes switching valves, an HP-PLOT Q column (30 m length x 0.53 mm i.d.; oven temperature program of 45°C for 6 min to 220 °C at 20°C/min) to separate fixed gases and methane from the remaining alkanes, as well as an HP-PLOT MoleSieve 5A (30 m length x 0.32 mm i.d.; oven temperature program of 50°C for 10 min to 120 °C at 10°C/min) for separation of fixed gases and methane. Helium was used as the carrier gas. After elution of methane, the second column is bypassed. Ethane and higher alkanes elute from the first column to a flame ionization detector. Methane and fixed gases are quantified on a thermal conductivity detector. Relative standard deviations were below 0.8 %.

Stable carbon and hydrogen isotope analyses of the individual compounds in the gas samples were performed repeatedly by gas chromatography-isotope ratio mass spectrometry using a GC/C III interface connected through a combustion reactor to a Finnigan MAT Delta plus XL mass

spectrometer. An Agilent 6890 series gas chromatograph equipped with a split-splitless capillary injector and a fused silica DB-1 capillary column (65 m length x 0.25 mm i.d.) was used. Gas samples were analyzed using a 100:1 split ratio injection system and the oven temperature program was 35°C (hold for 2 min) to 300 °C at 4 °C/min rate and was then held constant for 30 minutes. Helium was used as the carrier gas. Individual compounds separated by GC were pyrolyzed to CO₂ and H₂ at 850 °C and 1440 °C, respectively. Isotope compositions of N₂ were determined by the same mass spectrometer after separation from the gas samples in a glass vacuum line using a trap held at -196 °C. The ¹³C/¹²C, ¹⁵N/¹⁴N and D/H ratios are reported in “δ” notation. δ¹³C, δ¹⁵N and δD refer to V-PDB, N-SVEC and SMOW standards for carbon, nitrogen and hydrogen stable isotope measurements. Reproducibilities for δ¹³C, δ¹⁵N and δD values were better than ± 0.3 ‰, ± 0.5 ‰ and ± 5 ‰ for the dataset.

The API gravity of each oil sample was measured by ASTM D-4052 procedure (ASTM, 2011) through a SBS-3500 specific gravity meter that uses oscillating U-tube technology. The sample is filled into a container to measure the frequency of oscillation, from which the density value is calculated. Also, sulfur percentages were determined by ASTM D-4294 method (ASTM, 2010) using an energy-dispersive X-ray Panalytical spectrometer (Axios model) equipped with a digital signal processor and dual multi-channel analyser. Finally, we applied inductively-coupled plasma atomic emission spectroscopy through a Perkin-Elmer Optima 3000 spectrometer equipped with an automatic sampler for the quantitative determination of V and Ni contents (ASTM D-5708; ASTM, 2005).

4. Results and discussion

4.1. Chemical and isotopic composition of gases

Molecular concentrations of methane, ethane, propane, *n*-butane, iso-butane, *n*-pentane, iso-pentane, nitrogen, helium and argon are shown in Table 1. Methane was the main component in the gas samples, ranging between 27.97 and 75.5 % (average 62.9 %).

Table

1

Table 1

Average aliphatic hydrocarbons (<C₅), argon, and nitrogen from W-01 to W-11 gas samples.

	W-01	W-02	W-03	W-04	W-05	W-06	W-07	W-08	W-09	W-10	W-11
Methane	75.00	67.10	75.50	75.00	62.39	59.76	48.77	27.97	66.98	60.20	73.12
Ethane	13.40	14.40	13.65	13.55	15.77	18.89	16.24	22.55	16.17	16.90	14.82
Propane	7.81	10.93	6.90	7.43	11.99	12.22	16.57	26.12	10.23	13.10	7.68
Isobutane	0.72	1.17	0.83	0.59	1.85	1.43	2.23	3.27	1.21	1.69	0.85
n-butane	1.74	3.35	1.75	2.14	4.12	3.99	7.85	10.76	2.99	4.50	1.77
Isopentane	0.35	0.90	0.40	0.41	1.17	0.79	2.25	2.43	0.67	1.06	0.45
n-pentane	0.42	0.78	0.55	0.42	1.44	1.33	2.76	2.88	0.76	1.22	0.50
Nitrogen	0.27	0.27	0.27	0.29	0.27	0.29	0.27	0.26	0.29	0.97	0.91
Argon	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.02	0.002

The stable carbon isotopic compositions of the C₁-C₄ hydrocarbon compounds and hydrogen signature of methane for the gaseous samples are shown in Table 2. According to Schoell (1980), the $\delta^{13}\text{C}$ and δD ranges for methane in all samples indicate a thermogenic origin for this gas (Figure 3a). Likewise, based on a Bernard diagram (Figure 3b), $\delta^{13}\text{C}_1$ values over -55 ‰, uniform $\delta^{13}\text{C}_2$ data, and wetness values over 0.23, the natural gas samples could be classified as thermogenic gases generated from mature, sapropelic La Luna source rocks (Berner and Faber, 1988; Prinzhofer and Huc, 1995). Also, assuming that the $\delta^{13}\text{C}$ value of the La Luna kerogen is approximately -26 ‰ (Zumberge, 1984; Clayton and Bjorøy, 1994), and using the model proposed by Chung et al. (1988), the “natural gas plots” (r^2 values above 0.99; see Figure 3c) confirm a non-existent or scarce bacteriogenic methane fraction commingled with thermogenic methane (Rangel et al., 2003).

Table 2:

$\delta^{15}\text{N}$ data (‰), δD (‰) in methane and $\delta^{13}\text{C}$ values (‰) in methane, ethane, propane, isobutane and n-butane for the gas samples.

	$\delta^{13}\text{C}_1$	$\delta^{13}\text{C}_2$	$\delta^{13}\text{C}_3$	$\delta^{13}\text{iC}_4$		$\delta^{13}\text{nC}_4$	δD	$\delta^{15}\text{N}$
W-01	-51.31	-39.93	-32.78	-29.77		-28.98	-209	-9.33
W-02	-49.02	-39.09	-32.63	-29.93		-28.96	-206	-9.29
W-03	-51.12	-40.58	-32.92	-29.60		-29.23	-209	-9.26
W-04	-50.90	-39.82	-32.61	-29.84		-29.16	-208	-9.10
W-05	-51.13	-39.64	-32.80	-29.72		-29.28	-209	-9.36
W-06	-50.91	-39.31	-32.75	-29.93		-29.21	-208	-9.14
W-07	-51.06	-39.16	-32.55	-29.73		-29.33	-208	-9.30
W-08	-51.56	-39.73	-32.83	-29.84		-29.46	-209	-9.40
W-09	-50.12	-38.52	-32.25	-29.66		-28.91	-207	-9.19
W-10	-48.43	-37.12	-31.44	-29.71		-28.46	-204	-8.84
W-11	-48.00	-36.55	-31.27	-29.67		-27.99	-204	-8.89




Figure 3

Fig. 3. a) and b), respectively, relationships of δD to $\delta^{13}\text{C}_1$ and of $\delta^{13}\text{C}_1$ to $\text{C}_1/(\text{C}_2 + \text{C}_3)$ for the studied gases; c) $\delta^{13}\text{C}$

vs. the inverse C number of individual C₁–C₄ *n*-alkanes.

The molar concentrations of the C₁–C₄ alkanes indicate similar patterns, with two remarkable exceptions. Firstly, the W-08 sample showed an abnormally low value of methane content; in fact, in this well, a laboratory pressure test revealed a cylinder leak in coherence with chemical analysis, given that a notably lower concentration of methane, the major component, alters the relative concentration distribution pattern in the other organic compounds. An anomalous low concentration of methane was also noticed in W-07, but further pressure checking or geochemical analyses did not suggest anomalies in this well. However, as the gas-hydrocarbon sampling procedure is affected by production methods, pressure variation, and GOR (gas-oil ratio), changes in the pattern of this last well might be of circumstantial importance (Van Orsdal, 1990). These results confirm that it is not possible to obtain information about structural compartments using only the alkane (>C₁) compositions of the gas phase.

Nitrogen concentrations in gas samples at the well-head sampling sites provided valuable information. Gases from wells 10 and 11 showed molar nitrogen percentages (above 0.90 %) four times higher than those from the other wells (values between 0.26 and 0.29 %). These significant differences indicate that lateral composition is not equilibrated across the study reservoir. This chemically noteworthy step-like break could be indicative of the effect of geological barriers in molecular diffusion or convection, as discussed below.

Nitrogen is associated with the generation of high temperatures (Mingram et al., 2003), thus elevated concentrations of this gas are characteristic of a late hydrocarbon-pulse to the compartment without subsequent mixing with the rest of the study reservoir by diffusion or convection (Littke et al., 1995).

Assuming a high average N₂/Ar ratio above 200 and δ¹⁵N values about –9 ‰ in our case (Jenden et al., 1988; Zhu et al., 2000), the subsequent minimal or no contributions of magmatic, metamorphic or atmospheric N₂ lead us to infer that increasing concentrations of N₂ in the southern compartment are associated with increasing maturity levels in the source rock toward the south. In fact, these results are

consistent with other data discussed in this study.

Methane showed a typically lighter carbon isotopic ratio on average than average $\delta^{13}\text{C}_2$, whereas the $\delta^{13}\text{C}$ patterns of C_1 - C_4 alkanes ($\text{C}_1 < \text{C}_2 < \text{C}_3 < \text{C}_4$) were consistent with a regular Rayleigh “fractionation” model (Clayton, 1991). In particular, the isotopic compositions of methane and ethane from W-10 and W-11 presented differences of 2-3 ‰ with respect to the rest of the samples (see Table 2), thereby suggesting distinctive features. The gas correlation plot in Figure 4 again shows a clear compositional change in the gases from the southern area when compared with mean values of the sample set from the central and northern parts, which had low $\delta^{13}\text{C}$ standard deviations (< 1 ‰). In contrast, $\delta^{13}\text{C}$ for methane, ethane, propane, and *n*-butane in W-10 and W-11 were within experimental error range (0.2-0.3 ‰), thus indicating that these two samples belong to the same compartment. These results support the abovementioned chemical composition analyses (N_2) and indicate the presence of non-communicating reservoir compartments.




Fig. 4. $\delta^{13}\text{C}$ values (‰) in methane, ethane and propane from W-10, W-11, and the mean of the remaining sampled gases.

According to Lorant et al. (1998), isotopic data were also useful to hypothesize changes in the maturity levels of the samples. In fact, although all the gas samples analyzed fit within the oil window (see Figure 5a showing a diagram with vitrinite reflectances), heavier $\delta^{13}\text{C}$ values for W-10 and W-11 maybe related to increasing maturity levels in the source rock (Chung and Sackett, 1979), as reflected by increasing nitrogen values in the southern part of the reservoir. Secondary cracking or biodegradation in the reservoir compartments under study can be ruled out using the scheme of Prinzhofer et al. (2000), as indicated by the low C_2/C_3 ratios observed in all the gas samples (Figure 5b).




Fig. 5. a) and b), respectively, plots ($\delta^{13}\text{C}_2 - \delta^{13}\text{C}_3$ vs. C_2/C_3) and ($\text{C}_2/i\text{C}_4$ against C_2/C_3) for the gas samples.

Finally, natural gas data are described using star-diagrams applied to gas (also named Gastar diagrams by Prinzhofer and colleagues in 2000) to study the main processes (such as maturity, efficiency of accumulation or segregative migration) affecting the chemical and isotopic signatures of the samples. Eleven normalized geochemical parameters (C_1/C_2 , C_2/C_3 , $i\text{C}_4/n\text{C}_4$, $\delta^{13}\text{C}_1$, $\delta^{13}\text{C}_2$, $\delta^{13}\text{C}_3$, $\delta^{13}i\text{C}_4$, $\delta^{13}n\text{C}_4$, $\delta^{13}\text{C}_3 - \delta^{13}\text{C}_2$, $\delta^{13}n\text{C}_4 - \delta^{13}i\text{C}_4$, and $\delta^{13}\text{C}_2 - \delta^{13}\text{C}_1$) were discussed through a method reported by Prinzhofer et

al. (2000). All of these parameters were positively correlated with maturity, three (the isotopic ratios differences) were linked to the efficiency of accumulation, and another three (those involving methane) were strongly affected by migration. Figure 6 shows the aforementioned data plotted in three star-diagrams. As expected, the W-10 and W-11 samples can be grouped, both registering the greatest degree of maturity (Figure 6a). As already proposed, samples W-10 and W-11 represent a separate gas condensate reservoir compartment, which could be explained by differential entrapment of oil and gas.

In addition, the geochemical differences of gaseous samples from the northern and central parts of the study reservoir support the notion of at least two oil compartments. Figure 6b displays a series of natural gases (W-05 to W-09) with minor maturities, segregative fractionations and efficiencies of accumulation from the central part of the reservoir, as well as slight variations in these three processes. Although the differences between these five samples are very close to analytical accuracy, a reasonable connectivity in the central part of the reservoir could be inferred from the C₁-C₄ hydrocarbon analyses, which reveal a somewhat inefficient convection mechanism for mixing the fluids in a large reservoir (approximately 40 km² of areal extension). Consequently, it is not possible to conclude complete fluid homogenization.




Fig. 6. a), b) and c), three normalized Gostar diagrams for the groups of gas samples coming from the studied reservoir. The scales are the same for all 11 axes of each diagram.

In contrast, Figure 6c corresponds to the rest of the samples (gases W-01 to W-04) from the northern part of the reservoir and shows different behaviours; firstly, enhanced accumulation is observed, excluding W-01. Results for the W-02 sample showed compositional uniqueness; however, this feature is not enough to indicate a new single reservoir compartment or sub-compartment. Therefore, we cannot preclude that the hydrocarbon heterogeneities identified in Gostar diagrams are indicators of baffles and barriers to diffusion within the north-central part of the reservoir.

4.2. Petroleum bulk data heterogeneities

Results obtained for API gravity and sulfur content (Table 3) suggest intra-reservoir bulk lateral heterogeneity.

Table 3
Bulk geochemical data for the liquid samples.

	API	S	V	Ni	V/Ni
W-01	26.0	2.5	390	54	7.2
W-02	25.2	2.4	367	50	7.3
W-03	25.3	2.4	372	53	7.0
W-04	25.4	2.3	364	47	7.7
W-05	27.1	2.2	349	43	8.1
W-06	26.9	2.3	351	47	7.4
W-07	27.0	2.2	335	43	7.8
W-08	27.6	2.0	350	39	8.9
W-09	28.3	2.1	332	43	7.7
W-10	56.9	e	e	e	e
W-11	56.4	e	e	e	e
W-14	30.3	1.8	313	44	7.1

Note: V and Ni contents in ppm, sulfur ones in wt. %.

In this regard, the increasing API values of crude oils from the northeast to southwest may be attributable to the southward increase in maturity. Also, increasing maturity values from north to south may explain the southward decrease in the sulfur content of oils, which is possibly associated with lower concentrations of polar compounds (Peters et al., 2005). The V/Ni ratios above 7 (see Table 3) and sulfur content over 1% of the oils studied also pinpoint a carbonate source rock (La Luna Formation) deposited in an anoxic environment (Galarraga et al., 2008).

Consequently, this particular reservoir shows clear compositional trends from the northern to the southern part of the study area. The reservoir has a poor permeability (around 19 mD) and it is significantly long (11 km) and thin (16 m). These data can imply difficulties to obtain homogeneity of fluids (England et al., 1995). Computational simulation and/or detailed engineering monitoring should be used to examine possible stratigraphic transmissibility barriers in order to explain the fluid heterogeneities. On the basis of the data presented above, we propose that this area was filled from the

south, meaning that the source rock was in the southern part of the study area. Less mature oils accumulated in the north and successive petroleum charges filled the southern sands.

5. Conclusions

The isotopic and compositional signatures of individual C₁-C₄ aliphatic hydrocarbon compounds and nitrogen concentrations in gases presented a marked step-break variation between W-10 and W-11 and the rest of the gaseous samples toward the north. Specifically, nitrogen concentrations showed a notable step-break between these two populations, thereby suggesting a sealing character associated with a secondary fault identified in the geological section.

On the basis of the gas analyses, the presence of structural barriers in the central and northern parts of the reservoir is possible, since certain analytical trends do not rule out a scenario of additional single compartments or sub-compartments. Reservoir simulation or detailed engineering monitoring is required to locate additional transmissibility barriers.

The sulfur contents and specific gravities of the oils sampled corroborated a north-south compositional trend within the anticline structure and adjacent compartments. Based on the integration of all the geochemical parameters available, we propose that these reservoirs were filled from the south.

Acknowledgments

Authors are grateful to the company Perenco for accessing to the samples. We also thank the two anonymous reviewers for their comments which helped to improve the original version of this work

References

1. ASTM, 2005. ASTM D-5708 Standard for Determination of Nickel, Vanadium, and Iron in Crude Oils and Residual Fuels by Inductively Coupled Plasma (ICP) Atomic Emission Spectrometry: Annual Book of the American Society For Testing Materials Standards. West Conshohocken, ASTM International.
2. ASTM, 2010. ASTM D-4294 Standard for Sulfur in Petroleum and Petroleum Products by

Energy Dispersive X-ray Fluorescence Spectrometry: Annual Book of the American Society For Testing Materials Standards. West Conshohocken: ASTM International. ASTM, 2011. ASTM D-4052 Standard for Density, Relative Density, and API Gravity of Liquids by Digital Density Meter: Annual Book of the American Society For Testing Materials Standards. West Conshohocken: ASTM International.

3. Ballentine, C.J., Sherwood Lollar, B., 2002. Regional groundwater focusing of nitrogen and noble gases into the Hugoton-Panhandle giant gas field, USA. *Geochimica et Cosmochimica Acta*, 66, 2483-2497.
4. Bazan, L.W., 1998. The allocation of gas well production data using isotope analysis. SPE 40032, SPE Gas Technology Symposium, 15-18 March, Calgary, Alberta.
5. Beeunas, M.A., Baskin D.K., Jurgens, J., Dincau, A., Schoell, M., 1996. Application of gas geochemistry for reservoir continuity assesment, Gulf of Mexico. Abstract, AAPG-EAGE Research Symposium on "Compartmentalized Reservoirs: their detection, characterization and management", Houston.
6. Beeunas, M.A., Baskin, D.K., Schoell, M., 1999. Application of gas geochemistry for reservoir continuity assessment and identification of fault seal breakdown, South Marsh Island 61, Gulf of Mexico. Abstract, AAPG Hedberg Research Conference "Natural Gas Formation and Occurrence" June 6-10, Durango, Colorado.
7. Berner, U., Faber, E., 1988. Maturity related mixing model for methane, ethane and propane, based on carbon isotopes. *Organic Geochemistry*, 13, 67-72.
8. Chung H.M., Sackett, W.M., 1979. Use of stable carbon isotope compositions of pyrolytically derived methane as maturity indices for carbonaceous material: *Geochimica et Cosmochimica Acta*, 43, 1979- 1988.
9. Chung, H.M., Gormly, J.R., Squieres, R.M., 1988. Origin and of gaseous hydrocarbons in subsurface environment: theoretical considerations of carbon isotope distribution. *Chemical Geology*, 71, 97-103.

10. Clayton, C., 1991. Carbon isotope fractionation during natural gas generation from kerogen. *Marine and Petroleum Geology*, 8, 232-240.
11. Clayton, C., Bjoroy, M., 1994. Effect of maturity on $^{13}\text{C}/^{12}\text{C}$ ratios of individual compounds in North sea oils. *Organic Geochemistry*, 21, 737-750.
12. Clayton, J., 1997. Geochemistry of coalbed gas: a review. *International Journal of Coal Geology*, 35, 159-173.
13. England, W.A., Muggeridge, A., Clifford, P., Tang, Z., 1995. Modelling density-driven mixing rates in petroleum reservoirs on geological time-scales, with application to the detection of barriers in the Forties Field (UKCS), in: Cubitt J. & England W. (Eds.), *The Geochemistry of Reservoirs*, Special Publication No. 86, The Geological Society, Cambridge, pp. 185-201.
14. England, W.A., 2007. Reservoir Geochemistry: a reservoir engineering perspective. *Journal of Petroleum Science and Engineering*, 58, 344-354.
15. Erlich, R.N., Macsotay, O., Nederbragt, A.J., Antonieta Lorente, M., 2000. Birth and death of the Late Cretaceous “La Luna Sea”, and origin of the Tres Esquinas phosphorites. *Journal of South American Earth Sciences*, 13, 21-45.
16. Galarraga F, Reategui K, Martínez A, Martínez M, Llamas JF, Márquez G, 2008. V/Ni ratio as a parameter in palaeoenvironmental characterisation of nonmature medium-crude oils from several Latin American basins. *Journal of Petroleum Science and Engineering*, 61, 9-14.
17. Gold, T., Held, M., 1987. Helium-nitrogen-methane systematics in natural gases of Texas and Kansas. *Journal of Petroleum Geology*, 10, 415-424.
18. González de Juana, C., Iturralde de Arozena, J., Picard, X., 1980. *Geology of Venezuela and the petroliferous Venezuelan basins*, Vol. I & II, Fonvises Editions, Caracas.
19. Holba, A.G., Dzou, L.I.P., Franks, S.G., May, S.J., Lenney T., 1996. Reservoir geochemistry of

- South Pass 61 Field, Gulf of Mexico: Compositional heterogeneities reflecting filling history and biodegradation. *Organic Geochemistry*, 24, 1179-1198.
20. Hwang, R.J., Ahmed A.S., Moldowan, J.M., 1994. Oil composition variation and reservoir continuity: Unity Field, Sudan. *Organic Geochemistry*, 21, 171-188.
21. Jenden, P.D., Kaplan, I.R., Poreda, R.J., Craig, H., 1988. Origin of nitrogen rich gases in the Californian Great Valley: evidence from helium, carbon and nitrogen isotope ratios. *Geochimica et Cosmochimica Acta*, 52, 851-861.
22. Kaufman, R.L., Ahmed, A.S., Elsinger R.L., 1990. Gas chromatography as a development and production tool for fingerprinting oils from individual reservoirs: Applications in the Gulf of Mexico, in: *Proceedings of the ninth GCS-SEPM Annual Research Conference*, GCS-SEPM Foundation, Austin, pp. 263-282.
23. Kotarba, M.J., Nagao, K., 2008. Composition and origin of natural gases accumulated in the Polish and Ukrainian parts of the Carpathian region: gaseous hydrocarbons, noble gases, carbon dioxide and nitrogen. *Chemical Geology*, 255, 426-438.
24. Krooss, B.M., Leythaeuser, D., Lillack, H., 1993. Nitrogen-rich natural gases: qualitative and quantitative aspects of natural gas accumulation in reservoirs. *Erdöl und Hohle-Erdgas-Petrochemie*, 46, 271-276.
25. Larter, S.R., Aplin, A.C., 1994. Reservoir geochemistry: Methods, application and opportunities, in: England W.A. & Cubitt J. (Eds.), *The Geochemistry of Reservoirs*, Special Publication, No. 86, The Geological Society, Cambridge, pp. 5-32.
- 26.
27. Levaché, D., Montel, F., Walgenwitz, F., 2000. Deep offshore fluid evaluation/connectivity study. SPE 62904. SPE Annual Conference and Exhibition, 1-4 October, Dallas.

28. Littke, R., Krooss, B., Frielingsdorf, J., Muller, B., Schwochau, K., 1993. Generation of nitrogen and methane from carboniferous coals, in: Abstract Book of 16th International Meeting of Organic Geochemistry, The European Association of Organic Geochemists, Stavanger, pp. 43-50.
29. Littke, R., Krooss, B., Idize, E., Frielingsdorf, J., 1995. Molecular nitrogen in natural gas accumulations-generation from sedimentary organic matter at high temperatures. AAPG Bulletin, 79, 410-430.
30. Lorant, F.A., Prinzhofer, A.A., Behar, F., Huc, A.Y., 1998. Carbon isotopic and molecular constraints on the formation and the expulsion of thermogenic hydrocarbon gases. Chemical Geology, 147, 249- 264.
31. Milkov, A.V., Goebel E., Dzou, L., Fisher, D.A., Kutch, A., McCaslin, N., Bergman, D.F., 2007. Compartmentalization and time-lapse geochemical reservoir surveillance of the Horn Mountain oilfield, deep-water Gulf of Mexico. AAPG Bulletin, 91, 847-876.
32. Mingram, B., Hoth, P., Harlov, D.E., 2003. Nitrogen potential of Namurian shales in the North German Basin. Journal of Geochemical Exploration, 78/79, 405-408.
33. Parnaud, F., Gou, Y., Pascual, J.C., Capello, M.A., Truskowski, I., Passalacqua, H., 1995. Stratigraphic synthesis of western Venezuela, in: Tankard, A.J., Suárez, R., Welsink H.J. (Eds.), Petroleum Basins of South America, AAPG Memoir 62, The American Association of Petroleum Geologists, Tulsa, pp. 667-679.

34. Permanyer, A., Doufi, L., Lahcini, A., Lamontagne, J., Kister K., 2002. FTIR and SWF spectroscopy applied to reservoir compartmentalization: A comparative study with gas chromatography fingerprints results. *Fuel*, 81, 861-866.
35. Peters, K.E., Walters, C.C., Moldowan, J.M., 2005. *The Biomarker Guide: Biomarkers and Isotopes in Petroleum Systems and Earth History*, 2nd edition. Cambridge University Press, Cambridge.
36. Prinzhofer, A.A., Huc, A.Y., 1995. Genetic and post-genetic molecular and isotopic fractionations in natural gases. *Chemical Geology*, 126, 281-290.
37. Prinzhofer, A.A., Mello, M.R., da Sila Freitas, L.C., Takaki, T., 2000. A new geochemical characterization of natural gas and its use in oil and gas evaluation, in: M.R. Mello, & B.J. Katz (Eds.), *Petroleum systems and South Atlantic margins*, AAPG Memoir 70, Tulsa, pp. 107-119.
38. Rangel A., Katz, B., Ramírez, V., Vaz dos Santos Neto E., 2003. Alternative interpretations as to the origin of the hydrocarbons of the Guajira Basin, Colombia. *Marine and Petroleum Geology*, 20, 129- 139.
39. Rein, E., Schulz, L.K., 2007. Applications of natural gas tracers in the detection of reservoir compartmentalization and production monitoring. *Journal of Petroleum Science and Engineering*, 58, 428-442.
40. Rohrback, B.G., Peters, K.E., Sweeney, R.E., Kaplan, I.R., 1983. Ammonia formation in laboratory simulated thermal maturation: implications related to the origin of nitrogen in natural gas, in: Bjorøy, M. (Ed.), *Advances in Organic Geochemistry*, 1981, J. Wiley & Sons, New York, pp. 819-823.
42. Schoell, W., 1980. The hydrogen and carbon isotopic composition of methane from natural gases of various origins. *Geochimica et Cosmochimica Acta*, 44, 649-661.

43. Schoell, W., Jenden, P., 1993. Isotope analyses of gases in gas field and gas storage, in: Paper SPE 26171, Society of Petroleum Engineers, Richardson (US), pp. 337-344.
44. Smalley, P.C., England W.E., El Rabaa, A.W.M., 1994. Reservoir compartmentalization assessed with fluid compositional data, in: SPE Reservoir Engineering, Society of Petroleum Engineers, Richardson (US), pp. 175-180.
45. Talukdar, S., Gallango, O., Chin-A-Lien, M., 1986. Generation and migration of hydrocarbon in the Maracaibo Basin, Venezuela: An integrated basin study. *Organic Geochemistry*, 10, 261-279.
46. Talukdar, S., Marcano, F., 1994. Petroleum system of the Maracaibo Basin, Venezuela, in: Magoon, L., Dow, W. (Eds.), *The Petroleum System-From Source to Trap*. AAPG Memoir, No. 60, The American Association of Petroleum Geologists, Tulsa, pp. 463-482.
47. Tocco, R., Alberdi, M., 1999. Geochemical study of Misoa Formation crude oils, Centro Lago Field, Lake Maracaibo, Western Venezuelan basin. *Marine and Petroleum Geology*, 16, 135-150.
48. Van Orsdal, F.G., 1990. Natural gas measurement, sampling, and analysis. *Journal of Petroleum Technology*, 42, 780-783.
49. Wavrek, D., Mosca, F., Chaouche, A., Jarvie, D., 2001. Geochemical insights to engineering problems: when does oil heterogeneity really indicate a compartment?, in: *Abstract Book of 20th International Meeting of Organic Geochemistry*, The European Association of Organic Geochemists, Nancy (France), pp. 75-79.
50. Zhu, Y., Shi, B., Fang, C., 2000. The isotopic compositions of molecular nitrogen: implications on their origins in natural gas accumulations. *Chemical Geology* 164, 321-330
51. Zumberge, J.E., 1984. Source rocks of the La Luna Formation (Upper Cretaceous) in the Middle Magdalena valley, Colombia, in: Palacas, J.G. (Ed.). *Petroleum Geochemistry and Source Rock*

Potential of carbonate Rocks, AAPG Studies in Geology, 18, 127-133.