

Freshwater Biology

ADDRESSING LONGITUDINAL CONNECTIVITY IN FRESHWATER SYSTEMATIC CONSERVATION PLANNING.

Journal:	<i>Freshwater Biology</i>
Manuscript ID:	FWB-S-Jul-08-0322.R1
Manuscript Type:	Special Issue
Date Submitted by the Author:	n/a
Complete List of Authors:	Hermoso, Virgilio; University of Huelva, Biología Ambiental y Salud Pública Linke, Simon; The Ecology Centre, School of Integrative Biology; eWater Prenda, Jose; Universidad de Huelva, Biología Ambiental y Salud Pública POSSINGHAM, Hugh; University of Queensland St Lucia - Australia, The Ecology Centre
Keywords:	Conservaion / biodiversity < Applied Issues, Rivers < Habitat, Community < Level of Organisation, Modelling / statistics < Process / Approach / Methods, Fish < Taxonomic Group / Assemblage



1
2
3 ADDRESSING LONGITUDINAL CONNECTIVITY IN FRESHWATER
4
5
6 SYSTEMATIC CONSERVATION PLANNING.
7

8 Hermoso, V.¹, Linke, S.², Prenda, J.¹ & Possingham, H. P.²
9

10
11
12 ¹ Departamento de Biología Ambiental y Salud Pública, Universidad de Huelva. Avda.
13 Andalucía s/n, 21071 Huelva, Spain.
14

15
16 ² The Ecology Centre, School of Integrative Biology, University of Queensland, St
17 Lucia, Australia.
18

19
20
21
22 Running head: Connectivity in freshwater conservation planning
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Freshwater conservation has received less attention than its terrestrial or marine counterparts, despite freshwater systems containing a considerable amount of the earth's biodiversity. Given the accelerated rate of change and intensive human use that freshwater ecosystems are submitted to, it is urgent to devote some attention to them. The application of existing conservation planning tools - such as Marxan - to riverine planning needs some adaptations to account for the special nature of these systems. Connectivity plays a key role in freshwater ecosystems – threats are mediated along river corridors and the health of the entire catchment influences. This needs to be considered in conservation planning approaches.
2. The probability of occurrence, obtained from MARS-GLM models, of nine native freshwater fish species in a Mediterranean river basin was used as features to develop spatial conservation priorities. The priorities accounted for complementarity and spatial design issues.
3. To deal with the connected nature of rivers, we modified Marxan's boundary length penalty, hence avoiding the selection of isolated planning units and forcing the inclusion of closer upstream areas. We introduced 'virtual boundaries' between non-headwater stream segments, and added distance-weighted penalties to the overall connectivity cost (CP) when stream segments upstream of the selected planning units are not selected.
4. This approach to prioritising connectivity rule is concordant with ecological theory, as it considers the natural and roughly exponential decay of upstream influences with distance. It allows accounting for the natural capacity of rivers to mitigate impacts when designing reserves. With a small emphasis on connectivity, Marxan prioritised natural corridors for longitudinal movements. In contrast, whole sub-basins were prioritised when connectivity was emphasized. Changing the relative emphasis on connectivity causes substantial changes in the spatial prioritisation; our conservation investment could move from one basin to another.
5. Our novel approach to dealing with directional connectivity enables managers in charge of freshwater systems to set ecologically meaningful spatial conservation priorities.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

KEYWORDS: Biodiversity, irreplaceability, MARS-GLM, Marxan, Mediterranean, native fish species.

Copy for Review

INTRODUCTION

Despite containing a considerable amount of the Earth's global biodiversity (Allan & Flecker, 1993), and being exposed to higher pressures and threats than adjacent terrestrial ecosystems (Malmqvist & Rundle, 2002; Nel *et al.*, 2007), freshwater ecosystems have received less attention by the conservation community (Abell, 2002). There has been little emphasis on declaring protected areas for the primary purpose of conserving freshwater ecosystems and biodiversity (Saunders, Meewing & Vincent, 2002). Rivers have generally been inadequately dealt with in most assessments of terrestrial ecosystems unless they were considered important for terrestrial biodiversity patterns and processes (Nel *et al.*, 2007). Indeed, one of the primary uses of rivers in terrestrial reserves is to define reserve boundaries. We urgently need better conceptual frameworks and tools for freshwater conservation planning.

The protection of all places which contribute to biodiversity conservation is impossible, since conservation usually competes with other human interests (Margules, Pressey & Williams, 2002). Representativeness (adequate representation of all the targeted biodiversity attributes) and persistence of biodiversity are two main goals in reserve design (Margules & Pressey, 2000). Once established, reserves should promote the long term survival of the biodiversity they contain by maintaining natural processes and viable populations and by mitigating at least some of the proximate threats to their biodiversity (Margules & Pressey, 2000; Margules *et al.*, 2002; Wilson *et al.*, 2005a). Before systematic conservation planning, the acquisition of land for reserves traditionally involved either the use of a subjective judgment of biodiversity value, or the use of other completely extraneous criteria to biodiversity conservation such as scenic value, wilderness quality and inaccessibility, low primary production potential or simply availability (Margules, Nicholls & Pressey, 1988; Pressey, Possingham & Margules, 1996; Sarkar, 1999). These approaches lead to *ad hoc* conservation strategies focused on areas easiest to reserve, sometimes with least need for urgent or immediate protection (Pressey, 1994; Knight, 1999; Pressey *et al.*, 2000). In general, these kinds of approaches have not realised the benefits to biodiversity that they could have, as (i) they do not address representativeness (Pressey & Tully, 1994) and (ii) often force misallocation of limited resources into areas containing relatively few diversity surrogates (see later). To overcome these pitfalls, an explicit framework for systematic conservation planning has emerged (Margules & Pressey, 2000) in the last two decades.

1
2
3 Systematic conservation planning work tries to attain biodiversity conservation goals by
4 identifying important areas where conservation efforts should be focused (priority areas
5 hereafter) to facilitate the effective use of the limited resources intended for
6
7 conservation issues (Knight *et al.*, 2007).
8
9

10
11 Given the accelerated rate of land use change and because biodiversity protection
12 competes with legitimate alternative human uses, methods for identifying priority areas
13 need to be explicit, efficient, cost-effective and flexible (Margules *et al.*, 2002;
14 Possingham *et al.*, 2006). While traditional conservation tools relied on crude scoring
15 approaches based on different criteria such as species richness, rarity value, naturalness,
16 or size (Williams *et al.*, 1996), modern conservation planning methods use
17 complementarity-based algorithms and proper problem definition. Complementarity is
18 defined as the gain in representativeness of biodiversity when a site is added to an
19 existing set of areas (Possingham, Ball & Andelman, 2000). Algorithms which
20 incorporate complementarity lead to a more efficient representation of biodiversity
21 features and better cost-effective solutions than ad-hoc (Pressey & Tully, 1994), scoring
22 or ranking strategies (Margules *et al.*, 2002; Pressey & Nicholls, 1989). After defining
23 clear objectives (such as biodiversity targets), they look for areas that add as many
24 under-represented surrogates (taxa or any other conservation feature) as possible to a
25 network of protected areas (Pressey, Possingham & Day, 1997), achieving the
26 efficiency goal by selecting as few areas as possible that together reach the
27 representativeness goal (Pressey & Nicholls, 1989). However, the identification of
28 single reserve solutions is a rigid strategy which gives no indication on the importance
29 of each area in terms of their potential to be replaced by other available areas in the
30 region (Pressey, Watts & Barret, 2004) and the value of unselected areas (Cabeza &
31 Moilanen, 2006). To include flexibility in systematic conservation planning,
32 quantitative conservation tools often incorporate measures of irreplaceability –
33 calculated by estimating the likelihood that an area will be required to meet a given set
34 of targets (Pressey *et al.*, 1994; Ferrier *et al.*, 2000).
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 Most conservation planning approaches have so far overlooked freshwater
55 biodiversity, because incorporating freshwater species and habitats adds several layers
56 of complexity to an already complicated task (Abell, 2002). Nevertheless, systematic
57 conservation planning studies specifically targeting freshwater ecosystems have started
58 to emerge (Nel *et al.*, 2007; Linke *et al.*, 2007; Moilanen, Leathwick & Elith, 2008).
59
60

1
2
3 These studies apply all the principles developed for terrestrial ecosystems, but recognize
4 the need for some refinements to consider the special characteristics of freshwater
5 ecosystems (Dunn, 2003). Freshwater conservation planning must deal with the
6 connected nature of rivers, which is a key factor for the structure and conservation of
7 freshwater biodiversity - even more important in this context than for terrestrial
8 ecosystems (Ward et al., 2002). Riverine systems are characterised by multiple
9 longitudinal, lateral and vertical boundaries (Ward & Stanford, 1989). All these
10 boundaries are more efficiently connected than in terrestrial ecosystems given the
11 density and viscosity of water and its directional flow (Weins, 2002). This strong
12 connectivity has important ecological consequences (Vannote et al., 1980) and clear
13 effects leading the spread of perturbations and threats along freshwater ecosystems.
14 Despite the crucial role that connectivity plays in riverine ecosystems, it has not
15 received the attention that it deserves (Pringle, 2001; Ward et al., 2002). Reserves
16 located in middle or lower watersheds often suffer the cumulative effects of hydrologic
17 alteration and pollution originated in both upstream and downstream, imposing high
18 threats to the conservation of its biodiversity.
19
20
21
22
23
24
25
26
27
28
29
30
31

32 Here we adapt Marxan - an extensively used tool in terrestrial and marine
33 systematic conservation planning - to the peculiarities of rivers, aiming to attain more
34 effective reserve design principles for freshwater systems. We account for the connected
35 nature of rivers by introducing a penalty for not conserving upstream areas which could
36 have impacts on downstream reserves. In this way we specifically address one of the
37 components of spatial riverine connectivity, such as longitudinal connectivity, in
38 freshwater systematic conservation planning. At the same time, we simulate the decay
39 in spatial influence along water courses to reflect the natural capacity of rivers to
40 mitigate impacts.
41
42
43
44
45
46
47
48
49
50

51 METHODS

52 *Study area*

53
54
55 The Guadiana River basin is located in the South-Western Iberian Peninsula
56 draining a total area of 67039 Km² to the Atlantic Ocean (Fig. 1). It features a typical
57 Mediterranean climate, with high intra and inter-annual discharge variation, with severe
58
59
60

1
2
3 floods and droughts (Gasith & Resh, 1999). Mean air temperature ranges from 13 to
4 18.1 °C, with a strong intra-annual variation in extreme temperatures. Mean annual
5 precipitation ranges from 350 to 1200 mm (with a mean of 450 mm).
6
7

8
9 Although the Guadiana basin is not an overpopulated area (28 hab/km²),
10 agricultural activities have deeply transformed the landscape during the last century.
11 Almost half of the basin (49.1%) is currently used for agriculture - 30.6% occupied with
12 intensive agriculture as irrigated lands and 18.5% occupied with extensive agriculture,
13 like olive groves or fruit trees. As a consequence, about 8.3 10⁹ m³ of water is retained
14 in 86 big reservoirs (>10⁶ m³) and more than 200 small ones (<10⁶ m³) for water supply.
15 This has resulted in the modification of natural riverine flow regimes and has
16 fragmented fish habitat in the basin. Other common human perturbations are channel
17 modifications due to river channelization and degradation and even complete depletion
18 of the riparian forest. About 3,150 Km² (5.2% of basin's area) are formal reserves and
19 subject to special management regimes, though most of them arose from *ad hoc* or
20 terrestrial planning.
21
22
23
24
25
26
27
28
29

30
31 Guadiana's freshwater fish fauna is especially important within the circum-
32 Mediterranean context. Its high species richness is only comparable to that found in the
33 Po River basin in northern Italy and the lower Orontes in southern Turkey (Smith &
34 Darwall, 2006). All of these river basins contain between 11 and 17 native fish species.
35
36
37

38 *Planning units*

39
40
41 While in terrestrial systematic conservation planning equal sized grid cells are
42 often used as planning units, subcatchments are a more appropriate option for
43 freshwater environments. This spatial approach accounts for the connected nature of
44 rivers and natural boundaries of areas of influence (Linke et al., 2007; Klein et al., *in*
45 *press*). We derived 2170 planning units from a 90 m digital elevation model (Jarvis et
46 al., 2006) through ARC Hydro (Maidment, 2002) within ArcGIS 9.1 (ESRI, 2002).
47
48
49
50
51

52 *Environmental and biological data*

53
54
55 Presence/absence of fish species was determined by electrofishing in 151 sites,
56 each in a different planning unit. These sampling sites were homogeneously distributed
57 through the whole basin, ensuring an adequate characterization of natural variations and
58 human perturbations in the basin. Sampling was conducted once at every site without
59
60

1
2
3 block-nets in stretches of 100 m (when possible). This protocol is recommended for
4 obtaining an accurate characterization of species' presence-absence in the same area
5 (Filipe et al., 2004). The sampling stretch was representative of all the habitats present
6 in the area, including pools and riffles where available. All fish were released once they
7 were identified to species level.
8
9
10
11

12
13 Habitat data was used to build a predictive model with which relate freshwater
14 fish community and habitat characteristics. Then this model could be used to infer
15 freshwater fish communities from habitat data in unsampled planning units. The set of
16 environmental variables listed in Table 1, included a combination of both natural and
17 disturbance descriptors, to model actual probabilities of occurrence. We used two
18 different spatial scales to characterize the environmental attributes of our planning units
19 to portray both local influences and catchment scale effects, (i) subcatchment and (ii)
20 catchment scale. The mean value of each environmental variable in every planning unit
21 and across the whole upstream catchment area were considered respectively. Only
22 remotely sensed data was used in both approaches, to enable complete predictive
23 coverage. All the variables were tested for normality and appropriately transformed
24 when necessary before analysis.
25
26
27
28
29
30
31
32
33
34

35 *Prediction of biological data*

36
37
38 Lack of complete survey coverage is a common problem in conservation planning
39 (Margules & Pressey, 2000; Van Teeffelen, Cabeza & Moilanen, 2006; Linke *et al.*,
40 2007). This is usually dealt with by building predictive models of the distribution of
41 conservation features throughout the landscape (Wilson et al. 2005b). A Multivariate
42 Adaptive Regression Splines (MARS) model developed on the 151 sampled planning
43 units was used to predict the probability of occurrence for each species in the unsampled
44 planning units. MARS is a method of flexible non-parametric regression modelling
45 (Elith & Leathwick, 2007). It is useful for modelling complex non-linear relationships
46 between response and explanatory variables with similar levels of complexity to that of
47 a Generalized Additive Model (GAM) (Hastie, 1991). MARS fits a nonlinear function
48 to the relationships between dependent and predictor variables by breaking the range of
49 each predictor into a subset of portions or "knots", and fitting linear relationships for
50 each of them (basis functions). MARS allows the slope of the fitted linear segments
51 between pairs of segments to vary while ensuring that the full fitted function is without
52
53
54
55
56
57
58
59
60

1
2
3 breaks or sudden steps (Elith & Leathwick, 2007). The predictive function is finally
4 composed of a series of connected straight line segments, rather than the smooth curve
5 of a GAM. Two interesting features made MARS models useful for the present work. 1)
6 It allows exploring interactions between predictors (Leathwick, Elith & Hastie, 2006),
7 and 2) is able to fit a multi-response model which can simultaneously relate variation in
8 the occurrence of all species to the environmental predictors in one analysis (Olden,
9 2003). Multi-response models facilitate the modelling of rare species occurrences,
10 which are important in conservation planning exercises.
11
12
13
14
15
16
17

18 The model was fitted using a code provided by Elith & Leathwick (2007) for the
19 MDA (Mixture and Flexible Discriminant Analysis) library within the free statistical
20 software R, Version 2.1.1 (R Development Core TEAM, 2004). The common function
21 provided in R for MARS uses least squares which works appropriately for data with
22 normally distributed errors. With binomial data this results in the range of predicted
23 values being expanded beyond the acceptable range [0-1] (Leathwick *et al.*, 2006). To
24 solve this problem the cited code fits a MARS model using the standard R code,
25 extracts the basis functions, and computes a Generalized Linear Model (GLM) which
26 uses the basis functions as predictors of each species' presence-absence. This procedure
27 constrained predictions between of occurrence probabilities to between zero and one.
28 We allowed first order interactions between predictors in the models, since previous
29 analysis showed a significant improvement in model performance when they were
30 included. For more statistical details see Leathwick *et al.* (2005).
31
32
33
34
35
36
37
38
39
40
41

42 Model performance for each species was assessed through both measures of
43 deviance and the area under the receiver operating characteristic curve (ROC) (Fielding
44 and Bell, 1997). The area under the ROC curve (AUC) was assessed through a k-fold
45 cross validation procedure (Hastie, Tibshirani & Friedman, 2001). The data was
46 randomly divided in 10 exclusive sub-sets and model performance was calculated by
47 successively removing each sub-set, re-fitting the model with the remaining data, and
48 predicting the omitted data. The average error when predicting occurrence in new sites
49 can then be calculated by averaging the AUC across each of the subsets (Leathwick *et*
50 *al.*, 2005). An AUC>0.6 is usually defined as acceptable model performance (Fielding
51 & Bell, 1997). Deviance complements AUC because it expresses the magnitude of the
52 deviations of the fitted values from the observations. Analogous to Elith & Leathwick
53 (2007) the full information given in the predictions (raw probabilities of occurrence)
54
55
56
57
58
59
60

1
2
3 were used in both the AUC and deviance analysis, rather than transforming this data
4
5 into presence-absence estimates with a threshold.
6

7
8 Prior to model construction a Principal Component Analysis was carried out on
9
10 the environmental data to extract a reduced number of independent predictors for
11
12 MARS models. The presence-absence of 10 species in the whole biological data set was
13
14 used for fitting the models (n=151 planning units).
15

16 *Reserve design*

17
18 The best reserve system, and irreplaceability for each planning unit, were
19
20 calculated using the simulated annealing selection algorithm (Possingham *et al.*, 2000)
21
22 within the Marxan software package (Ball & Possingham, 2000). Marxan aims to find
23
24 an optimal reserve network by minimizing an objective function where feature
25
26 penalties, spatial design and cost tradeoffs are considered (Equation 1).
27

$$28 \text{ Objective function} = \sum_{\text{planning units}} \text{Cost} + \text{SPF} \sum_{\text{features}} \text{Feature Penalty} + \text{CP} \sum \text{ConnectivityCost}$$

29
30
31 Equation 1
32

33
34 The mathematical objective in Marxan is therefore:

35
36 *Minimise: The cost of all the sites in the reserve system plus a penalty of each feature*
37
38 *that does not reach its conservation target plus the cost of absent connections weighted*
39
40 *by CP, the “connectivity parameter”.*
41

42
43 After creating a random initial reserve system, planning units are added or
44
45 discarded from the reserve system in an attempt to minimise the objective function
46
47 (equation 1). The final aim is to adequately represent a set of targets (species in our
48
49 case) by selecting as few planning units as possible. In Equation 1, cost represents the
50
51 cost of preserving each planning unit. Since we lacked objective estimates of the
52
53 economic cost for the preservation of each planning unit, we assumed a homogeneous
54
55 cost for all of them.

56
57 The feature penalty (FP) is a penalty for not fully representing all the features
58
59 (fish species in this case) in the final reserve solution at the targeted level. Marxan
60
considers features as objectives rather than constraints so the final solution might fail to
meet adequate conservation for a feature if the weighting for the feature penalty is set

1
2
3 too low. We set the weight of the feature penalties for unmet targets high (SPF=100), so
4 all species targets were met. Species occurrences within the planning units were
5 formulated as probability of occurrence (p). Similar to Wilson et al. (2005), we did not
6 transform the probabilities into presence/absence, but treated the targets as expectations.
7 Hereby, if a target was 10, this target is fulfilled both by ten presences of $p_i=1$ being
8 reserved, as well as 20 presences of $p_i=0.5$ being reserved (cf Game et al. 2008). We set
9 a general target of 10 planning units, which roughly equates to 70 km of habitat for each
10 species.
11
12
13
14
15
16

17
18 In terrestrial applications, the spatial design of the reserve is determined by a
19 boundary length penalty that forces reserves to be compact. We modified the concept of
20 the boundary length penalty in Marxan to account for the connected nature of rivers (see
21 Linke *et al.*, 2007; Possingham *et al.* 2005). While Klein et al (in press) only consider
22 adjacent subcatchments, we introduce ‘virtual boundaries’ between non-headwater
23 subcatchments by adding penalties to the overall connectivity cost when subcatchments
24 upstream of the selected planning units are not selected (Fig. 2). Hereby, the penalty for
25 each planning unit decreases by a factor proportional to the reciprocal of the distance
26 between the planning units. A planning unit that is 1 km away from the selected
27 subcatchment incurs a penalty of 1, while a planning unit 2 km away incurs a penalty of
28 0.5 (= 1/2). Hence, the importance of an upstream subcatchment decays over the
29 distance to the planning unit containing the targets (Fig. 2).
30
31
32
33
34
35
36
37
38
39

40 How much emphasis we place on upstream connectivity can be adjusted using the
41 connectivity penalty (CP). A CP of 0, means that a planning unit can be chosen for
42 biodiversity protection without any incurring penalties for not including upstream
43 subcatchments. We tested the sensitivity of the reserve design outcomes to different
44 levels of the CP in order to find a reasonable value that balances connectivity with total
45 area to reserve. Ten different CPs were used (0, 0.001, 0.05, 0.1, 0.3, 0.5, 0.7, 1, 2, and
46 3). Finally, irreplaceability was assessed as the frequency of selection of each planning
47 unit by running the algorithm 100 times.
48
49
50
51
52
53
54

55 56 57 58 RESULTS

59 60 *Predictive models construction and performance*

1
2
3 The environmental variables with the highest loading for the 9 first Principal
4 Components (PC) of the environmental PCA were selected as predictors for the MARS-
5 GLM model (Table 2). These 9 PCs accounted for more than 76% of the original
6 variance and ensured high tolerance values (variance in each predictor not explained by
7 the remaining) for the variables used as predictors, avoiding redundancies.
8
9

10
11
12
13 Presence/absence of 9 out of the 10 species was successfully modelled with at
14 AUC>0.6 and an average explained deviance of 29% (Table 2), comparable to previous
15 studies (Leathwick *et al.*, 2005). The model only failed fitting *Cobitis paludica* data,
16 which is a ubiquitous species with high prevalence values which probably led to a
17 random distribution at least in relation to the selected predictors. This model was then
18 used to predict the probability of occurrence of each species in the unsampled planning
19 units.
20
21
22
23
24
25
26
27
28

29 *Reserve design*

30
31 As we increased the connectivity penalty (CP), we observed an exponential decay
32 in the ratio of boundary length to area needed (Fig. 3). This relationship was used to
33 identify a compromise setting of CP (CP = 0.01) at which there was a significant gain
34 for the reserve configuration in boundary length terms while keeping the total reserve
35 area reasonably low. In this way we prioritised for the most cost-efficient clustered
36 reserve (considering area as a surrogate for reserve cost).
37
38
39
40
41
42

43 Most species were overrepresented in the network for CP = 0.01 (Mean target
44 representation for all the species at CP = 0.01 was 13.7, although some species such as
45 *Iberocypris alburnoides*, reached a representation of 23.0, while others such as
46 *Luciobarbus comizo* and *Salaria fluviatilis* just reached the targeted level, 10). Both
47 these species had the highest probabilities of occurrence at downstream reaches. By
48 increasing the emphasis on upstream connectivity representing these two species forces
49 the inclusion of more planning units than necessary for the remaining species, hence
50 raising their representation within the reserve. Moreover, there was a roughly linear
51 increase in the area of the reserve system with increasing target levels (Fig 4).
52
53
54
55
56
57
58
59

60 Clearly, a high CP increased hydrologic connectivity of selected planning units
(Fig. 5c), since only headwater or whole upstream sub-catchments were included in the

1
2
3 best solution when a high CP was used. In contrast, isolated stream reaches, some even
4 located in the main Guadiana River channel, were selected in the best solution when the
5 CP was set at 0 (Fig. 5 a). Irreplaceability values, or frequency of selection of each
6 planning unit in 100 runs, followed a similar pattern with high values for isolated
7 planning units when using no CP (Fig. 5d) and more clustered solutions with the highest
8 irreplaceability values when setting a CP at higher values (Fig. 5 e and f). At increasing
9 CP values whole sub-basins were included in the best solution. Using a medium CP
10 (0.01) strings of planning units got selected (Fig. 5 b and e) – analogous to movement
11 corridors. At the higher setting (CP=3), entire subcatchments were included (Fig. 5 c
12 and f). Interestingly, the sub-catchments selected in ‘corridor mode’ (at an intermediate
13 CP, moderate upstream connectivity) were not the same as the sub-catchments
14 prioritised with a very high emphasis on upstream protection. When needing to protect
15 the entire basin, the solutions changed to a smaller basin – the Chanza River basin.
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 DISCUSSION

31
32 Systematic conservation planning aims to select a set of areas to efficiently ensure
33 the long-term persistence of targets (Margules & Pressey, 2000). The simulated
34 annealing optimization algorithm within Marxan increases efficiency in systematic
35 conservation planning. It is widely used by managers and discussed in the scientific
36 literature (Wilson *et al.*, 2005b; Oetting, Knight & Knight, 2006; Carwardine *et al.*,
37 2007). However some adaptations were needed to account for the connected nature of
38 rivers in freshwater systematic conservation planning. This is the first application in
39 which planning for upstream protection has been realized in Marxan, using freshwater
40 fish communities as surrogates.
41
42
43
44
45
46
47
48

49 Spatial connectivity is an issue of major concern in systematic conservation
50 planning (Cabeza, 2003), especially in freshwater applications due to the connected
51 nature of this environment. Pringle (2001) refers to four main patterns which have
52 important implications for the location and management of freshwater reserves, such as
53 (i) deterioration of lower watersheds, (ii) deterioration and loss of riverine floodplains,
54 (iii) deterioration of irrigated lands and connecting surface waters and (iv) isolation of
55 upper watersheds. All of these threats are connectivity-related. Hence, the consideration
56 of connectivity and its importance in maintaining natural ecological processes and
57
58
59
60

1
2
3 biodiversity in freshwaters is a key for systematic conservation planning in these
4 systems (Fausch et al., 2004; Ward, Malard & Tockner, 2004). With this purpose, we
5 modified the static rule proposed by Linke *et al.* (2007) - which requires all the
6 upstream river stretches to be included in the reserve - going beyond the optimization of
7 size, shape and other spatial issues of traditional conservation planning practices and
8 providing Marxan with a practical approach to tackle longitudinal connectivity.
9
10 Although connectivity penalties have been shown to affect the reserve configuration and
11 extent (see Carwardine *et al.*, 2007; Klein *et al.*, *in press*), there is a general agreement
12 in the benefits of this approach (Cabeza *et al.*, 2004). However, as commented above,
13 issues with riverine connectivity go beyond longitudinal aspects, having at least two
14 additional dimensions - lateral and vertical (Wiens, 2002). These should be addressed in
15 future approaches to fully consider river connectivity and ecological processes.
16
17
18
19
20
21
22
23
24

25
26 Increasing the connectivity penalty showed interesting and unexpected effects on
27 reserve design: at a value of CP around 0.01, the best solutions tended to be corridors
28 down a few river valleys. More emphasis on upstream connectivity forced the
29 prioritization of whole sub-basins. Moreover, both solutions were not nested. When
30 selecting full catchments, the Chanza river basin was chosen, while in 'corridor mode'
31 (lower CP at 0.01), the main conservation corridor was in a catchment further north.
32 This is an unusual response of Marxan, contradicting the linear responses to increasing
33 spatial reserve exigencies when enhancing the importance of reserve clustering. Stewart,
34 Noyce & Possingham (2003) found a nested pattern with increasing representation,
35 however in our example the focal catchment differed when increasing the CP. The
36 reason for this is that Degebe River has a larger catchment overall. Thus, cost for
37 whole-of-basin protection favours the smaller Chanza basin. At lower CPs in which the
38 entire basin does not need to be protected to get a good solution, the longer stream
39 network provides a better solution as a corridor. This fact is important to acknowledge
40 in the planning stage, as planning for corridors with a low CP will result in a completely
41 different reserve configuration to the whole-of-basin planning.
42
43
44
45
46
47
48
49
50
51
52
53

54
55 By introducing decreasing virtual boundary penalties towards upstream
56 catchments, we also simulated the natural decay in the influence of river segments on
57 lower reaches. The decay we included ensures an appropriate weighting of the potential
58 effect of upstream disturbances by their distance to the targeted area. This resembles the
59 behavior of natural systems (Prenda & Gallardo-Mayenco, 1996; Wiens, 2002), and is a
60

1
2
3 vast improvement to the rigid rule proposed by Linke et al. (2007). The selection of all
4 the upstream catchment is often unrealistic when dealing with conservation of lowland
5 reaches. Our approach has important implications for future consideration of threats to
6 the conservation in freshwater conservation planning, since reserves could be selected
7 accounting for weighted distances to the main perturbations (such as centers of exotic
8 species populations or highly human perturbed environments). Dealing with natural
9 processes has been flagged as a key issue for an effective systematic conservation
10 planning in the present changing world (Pressey *et al.*, 2007). The connectivity rule
11 addresses it not only through the selection of longitudinally connected reserves which
12 mitigates the drawbacks of present freshwater reservation highlighted by Pringle (2001)
13 and Oetting *et al.* (2006), but also accounting for the natural capacity of rivers to
14 mitigate impacts with distance along longitudinal gradients.
15
16
17
18
19
20
21
22
23
24

25
26 A common problem in conservation planning is that species distribution data is
27 often incomplete. To fill the gaps, predicted species distributions can be used (Cabeza,
28 2003; Cabeza *et al.*, 2004; Linke *et al.*, 2007). Two different approaches have been
29 previously followed at this stage: using direct probability of occurrence or transforming
30 them into presence-absence data through arbitrary thresholds. The latter approach has
31 been tested and used in previous studies (see Polasky *et al.*, 2000; Wilson *et al.*, 2005b).
32 Despite this suppose a more risk-averse approach to conservation planning (Wilson et
33 al., 2005b), it also entails a net loss of information on species distribution data. The
34 threshold used in this transformation influences not only the predicted distribution area
35 for the conservation features, but also the outputs of conservation planning process and
36 has to be carefully set to ensure a suitable use (Wilson *et al.*, 2005b). Moreover, some
37 reserve selection methods based on presence-absence data may fail to consider
38 persistence of targets in reserve selection (Araújo & Williams, 2000; or Teeffelen *et al.*,
39 2006). We dealt with persistence more thoroughly by using present probabilities of
40 occurrence (see Cabeza *et al.* 2004), instead of potential probabilities of sites used in the
41 reference condition approach (see Linke et al. 2007). This probability of occurrence
42 indicates the likelihood with which a species is present in a planning unit considering
43 different species-dependent factors such as habitat quality requirements or vulnerability
44 to threats (Araújo & Williams, 2000). Instead of using just reference site distributions
45 (Linke *et al.* 2007), our present distribution models were built on the whole dataset -
46 using even sites with perturbed fish communities and including human-influenced
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 environmental variables as predictors. Potential distributions can be unrealistic, since
4 some species may have been pushed out to marginal areas within their original
5 distribution or displaced to new areas due to human impairment (Kouamélan *et al.*,
6 2003; Light & Marchetti, 2007). In this sense, Araújo, Williams & Fuller (2002)
7 showed how the probability of persistence increases if reserve selection algorithms
8 maximize the probability of the current occurrence instead of using a hypothetical niche
9 model. With actual distributions we focused the efforts on identifying the more suitable
10 areas for attaining persistence and optimizing the use of the scarce resources intended
11 for conservation purposes (Knight *et al.*, 2007) in conservation areas under current land-
12 use, at least if they remain stable.
13
14
15
16
17
18
19
20
21

22 An additional benefit of using direct probabilities of occurrence is the ability to
23 better portray the continuous nature of rivers (Vannote *et al.*, 1980) in the conservation
24 process. Since biological communities change gradually through natural upstream-
25 downstream gradients in rivers (Allan, 1995; Welborn, Skelly & Werner, 1996;
26 Clavero, Blanco-Garrido & Prenda, 2005) spatial connectivity is better addressed
27 through continuous probabilities rather than through presence-absence data. It also
28 allows the interpretation of the reserve in terms of a trade-off between river length and
29 probability of occurrence. In our approach we ensured selecting 70 river kilometers
30 where the species had a high certainty of occurrence ($p_i=1$) or larger habitat length at a
31 lower probability. This could then be related to the spatial needs for each species to
32 develop healthy populations and their probability of persistence. A feedback process,
33 where this kind of basic ecological information will guide reserve selection through
34 setting variable representation targets, would have clear beneficial effects for the
35 effectiveness of the conservation plan (Pressey *et al.*, 2007). However, as this
36 information is still lacking we have to trust the river habitat length included in the
37 present best reserve to be enough for preserving all the targeted species.
38
39
40
41
42
43
44
45
46
47
48
49
50

51 A whole basin approach has been followed in this study while previous efforts in
52 the same area were only focused on the Portuguese (Filipe *et al.*, 2004) or Spanish
53 (Hermoso *et al.*, *submitted*) portions of the basin while cross-boundary conservation
54 management seems a more effective practice. The most irreplaceable area and the best
55 solution when setting the CP at high values were especially focused on a single sub-
56 basin (Chanza River). This solution highlights the importance of tributaries for
57 conserving freshwater biodiversity (see Nel *et al.*, 2007). The same area had previously
58
59
60

1
2
3 been included in a set of priority areas for conservation through an alternative method
4 based on the reference condition approach (Hermoso *et al.*, *submitted*). However this is
5 the first time that a complementarity-based algorithm has been applied to this basin in
6 particular. Algorithms which incorporate complementarity ensure representativeness, in
7 addition to persistence, which is the other major goal in systematic conservation
8 planning, (Margules & Pressey, 2000; Margules *et al.*, 2002). This guaranties the
9 adequate representation of each species within the reserve, overcoming uncovered
10 deficiencies in other reserve selection methods based in scoring and ranking approaches
11 (Williams *et al.*, 1996; Margules *et al.*, 2002).
12
13
14
15
16
17
18
19

20 This study introduces substantial innovations to freshwater systematic
21 conservation planning. We have modified Marxan to specifically deal with longitudinal
22 connectivity in freshwater conservation through the inclusion of our virtual boundaries
23 between planning units. A flexible connectivity penalty allowed the consideration of the
24 natural decay by distance between stream reaches. At different penalty strengths
25 Marxan was able to identify either longitudinal corridors or whole sub-basins (at
26 increasing CPs). All of these advances contribute to improved realism when dealing
27 with freshwater conservation issues.
28
29
30
31
32
33
34
35
36
37

38 ACKNOWLEDGEMENTS

39
40 The fieldwork was supported by the Confederación Hidrográfica del Guadiana through
41 the project “*Diseño y Explotación de la Red de Control Biológico de la Cuenca del Río*
42 *Guadiana*” and the Ministry of Science and Technology (REN2002-03513, CGL2005-
43 02699). We thank Noam Levin for the help with GIS information, Jane Elith for the
44 statistical help with MARS-GLM models, and everyone at the Ecology Centre of the
45 University of Queensland for early comments and suggestion on this work. V. Hermoso
46 holds a pre-doctoral fellowship financed by the Spanish Ministry of Education (AP-
47 2004-1414). Simon Linke’s postdoctoral fellowship is funded by the eWater CRC.
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

- 1
2
3
4
5
6 Abell R. (2002) Conservation biology for the biodiversity crisis: a freshwater follow-up.
7 *Conservation Biology*, **16**, 1435-1437.
8
- 9 Allan J. D. & Flecker A. S. (1993) Biodiversity conservation in running waters.
10 *Bioscience*, **43**, 32-43.
11
- 12 Allan, J. D. (1995) *Stream ecology- structure and function of running waters*. Chapman
13 and Hall, London, UK.
14
- 15 Araújo M. B. & Williams P. H. (2000) Selecting areas for species persistence using
16 occurrence data. *Biological Conservation*, **96**, 331-345.
17
- 18 Araújo M. B., Williams, P. H. & Fuller R. J. (2002) Dynamics of extinction and the
19 selection of nature reserves. *Proceedings of the Royal Society of London, B*, **269**,
20 1970-1980.
21
- 22 Ball I. R. & Possingham H. P. (2000) Marxan version 1.8.3.
23 <http://www.ecology.uq.edu.au/Marxan.htm#1>
24
- 25 Cabeza M. & Moilanen A. (2006) Replacement cost: A practical measure of site value
26 for cost-effective reserve planning. *Biological Conservation*, **132**, 336-342.
27
- 28 Cabeza M. (2003) Habitat lost and connectivity of reserve networks in probability
29 approaches to reserve design. *Ecological Letters*, **6**, 665-672.
30
- 31 Cabeza M., Araújo M. B., Wilson R. J., Thomas C. D., Cowley M. R. J. & Moilanen A.
32 (2004) Combining probabilities of occurrence with spatial reserve design. *Journal of*
33 *Applied Ecology*, **41**, 252-262.
34
- 35 Carwardine J., Rochester W. A., Richardson K. S., Williams K. J., Pressey R. L. &
36 Possingham H. P. (2007) Conservation planning with irreplaceability: does the
37 method matter? *Biodiversity and Conservation*, **16**, 245-258.
38
- 39 Clavero, M., Blanco-Garrido, F. & Prenda, J. (2005) Fish-habitat relationship and fish
40 conservation in small coastal streams in southern Spain. *Aquatic Conservation-*
41 *Marine and Freshwater Ecosystems*, **15**, 415-426.
42
- 43 Cowling R.M., Pressey R.L., Rouget M. & Lombard A.T. (2003) A conservation plan
44 for a global biodiversity hotspot - the Cape Floristic Region, South Africa.
45 *Biological Conservation*, **112**, 191-216.
46
- 47 Dunn H. (2003) Can conservation assessment criteria developed for terrestrial systems
48 be applied to riverine systems? *Aquatic Ecosystem Health and Management*, **6**, 81-
49 95.
50
- 51 Elith J. & Leathwick J. (2007) Predicting species distributions from museum and
52 herbarium records using multiresponse models fitted with multivariate adaptive
53 regression splines. *Diversity and Distributions*, **13**, 265-275.
54
- 55 ESRI. (2002) *ArcGIS*. Environmental Systems Research Institute, Redlands, CA.
56
- 57 Fausch K. D., Torgersen C. E., Baxter C. V. & Li H. W. (2004) Landscapes to
58 riverscapes: bridging the gap between research and conservation of stream fishes.
59 *Bioscience*, **52**, 483-498.
60

- 1
2
3 Ferrier S., Pressey R. L. & Barret T. W. (2000) A new predictor of the irreplaceability
4 of areas for achieving conservation goals, its application to real-world planning, and
5 a research agenda for further refinement. *Biological Conservation*, **93**, 303-325.
6
7 Fielding A. H. & Bell J. F. (1997) A review of methods for the assessment of prediction
8 errors in conservation presence/absence models. *Environmental Conservation*, **24**,
9 38-49.
10
11 Filipe A. F., Marques T. A., Seabra S., Tiago M. P., Ribeiro F., Moreira da Costa L.,
12 Cowx I. G. & Collares-Pereira M. J. (2004) Selection of priority areas for fish
13 conservation in Guadiana River basin, Iberian Peninsula. *Conservation Biology*, **18**,
14 189-200.
15
16 Game E. T., Watts M. E., Wooldridge S. and Possingham H. P. (2008) Planning for
17 persistence in marine reserves : a question of catastrophic importance. *Ecological*
18 *Applications*, **18**, 670-680.
19
20 Gasith A. & Resh V H. (1999) Streams in Mediterranean climate regions: abiotic
21 influences and biotic responses to predictable seasonal events. *Annual Review of*
22 *Ecology and Systematics*, **30**, 51-81.
23
24 Hastie T. (1991) *Generalized additive models*. In *Statistical Models* in S. Chambers, J.
25 M. and Hastie, T. J. (eds), pp: 249-308. Wadsworth and Brooks/Cole Advanced
26 Books and Software, California.
27
28 Hastie T., Tibshirani R. J. & Friedman J. H. (2001) *The elements of statistical learning:*
29 *data mining, inference and prediction*. Springer-Verlag, New York.
30
31 Hermoso V., Linke S. & Prenda J. (submitted). Identifying priority areas for
32 conservation of freshwater fish biodiversity in a Mediterranean basin with a high
33 degree of threaten endemics. *Hydrobiologia*.
34
35 Hijmans R.J., Cameron S.E., Parra J.L., Jones P.G. & Jarvis A. (2005) Very high
36 resolution interpolated climate surfaces for global land areas. *International Journal*
37 *of Climatology*, **25**, 1965-1978.
38
39 Howard P. C., Viskanac P., Davenport T. R. B., Kigenyi F. W., Baltzer M., Dickinson
40 C. J., Lwanga J. S., Matthews R. A. & Balmford A. (1998) Complementarity and
41 the use of indicator groups for reserve selection in Uganda. *Nature*, **394**, 472-475.
42
43 Jarvis A., Reuter H.I., Nelson A. & Guevara E. (2006) *Hole-filled seamless SRTM data*
44 *V3*, International Centre for Tropical Agriculture (CIAT), available from
45 <http://srtm.csi.cgiar.org>.
46
47 Klein C., Wilson K., Watts M., Stein J., Berry S., Carwardine J., Stafford Smith M.,
48 Mackey B. & Possingham H. (in press). Incorporating ecological and evolutionary
49 processes into continental scale conservation planning. *Ecological applications*.
50
51 Knight A. T., Smith R. J., Cowling R. M., Desmet P. G., Faith D. P., Ferrier S.,
52 Gelderblom C. M., Grantham H., Lombard A. T., Maze K., Nel J. L., Parrish J. D.,
53 Pence G. Q. K, Possingham H. P., Reyers B., Ruoget M., Roux D. & Wilson K. A.
54 (2007) Improving the key biodiversity areas approach for effective conservation
55 planning. *Bioscience*, **57**, 256-261.
56
57
58
59
60

- 1
2
3 Knight R. L. (1999) Private lands: the neglected geography. *Conservation Biology*, **13**,
4 223-224.
5
6 Kouamélan E. P., Teugels G. G., N'Douba V., Gooché Bi G. & Koné T. (2003) Fish
7 diversity and its relationships with environmental variables in a West African basin.
8 *Hydrobiologia*, **505**, 139-146.
9
10 Leathwick J. R., Elith J. & Hastie T. (2006) Comparative performance of generalized
11 additive models and multivariate adaptive regression splines for statistical modelling
12 of species distributions. *Ecological Modelling*, **190**, 188-196.
13
14 Leathwick J. R., Rowe D., Richardson J, Elith J. & Hastie T. (2005). Using multivariate
15 adaptive regression splines to predict the distribution of New Zealand's freshwater
16 diadromous fish. *Freshwater Biology*, **50**, 2034-2052.
17
18 Light T. & Marchetti M. P. (2007) Distinguishing between invasions and habitat
19 changes as drivers of diversity loss among California's freshwater fishes.
20 *Conservation Biology*, **21**, 434-446.
21
22 Linder Mayer D. B., Margules C. R. & Botkin, D. B. (2000) Indicators of biodiversity
23 for ecologically sustainable forest management. *Conservation Biology*, **4**, 941-950.
24
25 Linke S., Pressey R. L., Bailey R. C. & Norris R. H. (2007) Management options for
26 river conservation planning: condition and conservation re-visited. *Freshwater*
27 *Biology*, **52**, 918-938.
28
29 Maidment D. R. (2002) *Arc Hydro: GIS for Water Resources*. ESRI Press, Redlands,
30 CA.
31
32 Malmqvist B. & Rundle S. (2002) Threats to the running water ecosystems of the world.
33 *Environmental Conservation*, **29**, 134-153.
34
35 Margules C. R. & Pressey R. L. (2000) Systematic conservation planning. *Nature*, **405**,
36 243-253.
37
38 Margules C. R., Nicholls A. O. & Pressey R. L. (1988) Selecting networks of reserves
39 to maximize biological diversity. *Biological Conservation*, **43**, 63-76.
40
41 Margules C. R., Pressey R. L. & Williams, P. H. (2002) Representing biodiversity: data
42 and procedures for identifying priority areas for conservation. *Journal of*
43 *Biosciences*, **27**, 309-326.
44
45 Moilanen A., Leathwick J. & Elith J. (2008) A method for freshwater conservation
46 prioritization. *Freshwater Biology*, **53**, 577-592.
47
48 Nel J. L., Roux D. J., Maree G., Kleynhans C. J., Moolman J., Reyes B., Rouget M. &
49 Cowling R. M. (2007) Rivers in peril inside and outside protected areas: a
50 systematic approach to conservation assessment of river ecosystems. *Diversity and*
51 *Distributions*, **13**, 341-352.
52
53 Oetting J. B., Knight A. L. & Knight G. R. (2006) Systematic reserve design as a
54 dynamic process: F-TRAC and the Florida Forever Program. *Biological*
55 *Conservation*, **128**, 37-46.
56
57
58
59
60

- 1
2
3 Olden, J. D. (2003) A species-specific approach to modelling biological communities
4 and its potential for conservation. *Conservation Biology*, **17**, 854-863.
- 5
6 Poff N. L., Allan J. D., Bain M. B., Karr J. R., Prestegard K. L., Richter B. D., Sparks
7 R. E. & Stromberg J. C. (1997) The natural flow regime. A paradigm for river
8 conservation and restoration. *Bioscience*, **47**, 769-784.
- 9
10 Polasky S., Camm J. D., Slow A. R., Csuti B., White D. & Ding R. (2000) Choosing
11 reserve networks with incomplete species information. *Biological Conservation*, **94**,
12 1-10.
- 13
14 Possingham H. P., Ball I. R. and Andelman S. (2000) *Mathematical methods for*
15 *identifying representative reserve networks*. In: Ferson, S. and Burgman, M. (eds).
16 Quantitative methods for conservation biology. Springer-Verlag, New York, pp:
17 291-305.
- 18
19 Possingham H. P., Franklin J., Wilson K. & Regan T. J. (2005) The roles of spatial
20 heterogeneity and ecological processes in conservation planning. In: *Ecosystem*
21 *function in heterogeneous landscapes* (Eds. G.M. Lovett, C.G Jones, M.G. Turner
22 & K.C Weathers). Pp 389-406. Springer-Verlag, New York.
- 23
24 Possingham H. P., Wilson K. A, Andelman S. & Vynne C. H (2006) Protected Areas:
25 Goals, Limitations, and Design. In *Principles of Conservation Biology* (Eds. M. J.
26 Groom, G. K. Meffe, & C. R. Carroll) Pp 509-533. 3rd Edition. Sinauer Associates,
27 Inc., Sunderland MA.
- 28
29 Prenda, J. & Gallardo-Mayenco, A. (1996) Self-purification, temporal variability and
30 the macroinvertebrate community in small lowland Mediterranean streams receiving
31 crude domestic sewage effluents. *Archiv für Hydrobiologie*, **136**, 159-170.
- 32
33 Pressey R. L. & Nicholls A. O. (1989) Efficiency in conservation evaluation: scoring
34 versus iterative approaches. *Biological Conservation*, **50**, 199-218.
- 35
36 Pressey R. L. & Tully S. L. (1994) The cost of ad hoc reservation: a case study in New
37 South Wales. *Australian Journal of Ecology*, **19**, 375-384.
- 38
39 Pressey R. L. (1994) Ad hoc reservations: forward and backward steps in developing
40 representative reserve systems. *Conservation Biology*, **8**, 662-668.
- 41
42 Pressey R. L., Cabeza M., Watts M. E., Cowling R. M. & Wilson K. A. (2007)
43 Conservation planning in a changing world. *Trends in Ecology and Evolution*, **22**,
44 583-592.
- 45
46 Pressey R. L., Hager T. C., Ryan K. M., Schwarz J., Wall S., Ferrier S. & Creaser P. M.
47 (2000) Using abiotic data for conservation assessments over extensive regions:
48 quantitative methods applied across New South Wales, Australia. *Biological*
49 *Conservation*, **96**, 55-82.
- 50
51 Pressey R. L., Humphries C. J., Margues C. R., Vane-Wright, R. I. & Williams P. H.
52 (1993) Beyond opportunism: key principles for systematic reserve selection. *Trends*
53 *in Ecology and Evolution*, **8**, 124-128.
- 54
55
56
57
58
59
60

- 1
2
3 Pressey R. L., Possingham H. P. & Day J. R. (1997) Effectiveness of alternative
4 heuristic algorithms for identifying minimum requirements for conservation
5 reserves. *Biological Conservation*, **80**, 207-219.
6
7
8 Pressey R. L., Possingham H. P. & Margules C. R. 1996. Optimality in reserve selection
9 algorithms: when does it matter and how much? *Biological Conservation*, **76**, 259-
10 267.
11
12 Pressey R. L., Watts M. E. & Barret T. V. (2004) Is maximizing protection the same as
13 minimizing loss? Efficiency and retention as alternative measures of the
14 effectiveness of proposed reserves. *Ecological Letters*, **7**, 1035-1046.
15
16 Pressey R.L., Hager T. C., Ryan K. M., Schwarz J., Wall S., Ferrier S. & Creaser P. M.
17 (2000) Using abiotic data for conservation assessments over extensive regions:
18 quantitative methods applied across New South Wales, Australia. *Biological*
19 *Conservation*, **96**, 55-82.
20
21 Pringle C. M. (2001) Hydrologic connectivity and the management of biological
22 reserves: A global perspective. *Ecological Applications*, **11**, 981-998.
23
24 R Development Core Team. (2004) *A language and environment for statistical*
25 *computing*. R Foundation for Statistical Computing, Viena, Austria. ISBN 3-
26 900051-07-0, URL <http://www.R-project.org>.
27
28 Sarkar S. (1999) Wilderness preservation of biodiversity conservation-Keeping
29 divergent goals distinct. *Bioscience*, **49**, 405-412.
30
31 Saunders D. L., Meeuwig J. J., & Vincent C. J. (2002) Freshwater protected areas:
32 strategies for conservation. *Conservation Biology*, **16**, 30-41.
33
34 Smith K. G. & Darwall W. R. T. (Ed.), 2006. *The Status and Distribution of Freshwater*
35 *Fish Endemic to the Mediterranean Basin*. IUCN, Gland, Switzerland and
36 Cambridge, UK. v + 34 pp.
37
38 Stewart R. R., Noyce T. & Possingham H. P. (2003) Opportunity cost of ad hoc marine
39 reserve design decisions: an example from South Australia. *Marine Ecology*
40 *Progress Series*, **253**, 25-38.
41
42 Van Teeffelen A. J. A., Cabeza M. & Moilanen A. (2006) Connectivity, probabilities
43 and persistence: comparing reserve selection strategies. *Biodiversity and*
44 *Conservation*, **15**, 899-919.
45
46 Vannote R. L., Minshall G. W., Cummins K. W., Sedell J. R. & Cushing C. E. (1980)
47 River continuum concept. *Canadian Journal of Fisheries and Sciences*, **37**, 130-
48 137.
49
50 Ward J. V. & Stanford J. A. (1989) The four-dimensional nature of lotic systems.
51 *Journal of the North American Benthological Society*, **8**, 2-8.
52
53 Ward J. V., Malard F. & Tockner K. (2004) Landscape ecology: a framework for
54 integrating patterns and process in river corridors. *Landscape Ecology*, **17**
55 (Supplement 1), 35-45. Abell R. (2002) Conservation biology for the biodiversity
56 crisis: a freshwater follow-up. *Conservation Biology*, **16**, 1435-1437.
57
58
59
60

- 1
2
3 Ward J. V., Tockner K., Arscott D. B. & Claret C. (2002) Riverine landscape diversity.
4 *Freshwater Biology*, **47**, 517-539.
5
6 Weins J. A. (2002) Riverine landscapes: taking landscape ecology into the water.
7 *Freshwater Biology*, **47**, 501-515.
8
9 Welborn, G. A., Skelly, D. K. & Werner, E. E. (1996) Mechanisms creating community
10 structure across a freshwater habitat gradient. *Annual Review of Ecology and*
11 *Systematics*, **27**, 337-363.
12
13 Williams P. H., Humphries C. J., Araujo M. B., Lahti T., Lampinen R., Uotila P. &
14 Vane-Wright R. I. 1999. *Important plant areas of Europe: exploring consequence of*
15 *selection criteria; in Planta Europa*. Synge, H. and Akeroyd, J. (eds), London
16 Plantlife, pp: 103-109.
17
18 Williams P., Gibbons D., Margules C., Rebelo A., Humphries C. & Pressey R. (1996) A
19 comparison of richness hotspots, rarity hotspots and complementarity areas for
20 conserving diversity of British birds. *Conservation Biology*, **10**, 155-174.
21
22 Wilson K. A., Westphal M. I., Possingham H. P. & Elith J. (2005b) Sensitivity of
23 conservation planning to different approaches to using predicted species distribution
24 data. *Biological Conservation*, **122**, 99-112.
25
26 Wilson K., Pressey R. L., Newton A., Burgman M., Possingham H & Weston C.
27 (2005a) Measuring and incorporating vulnerability into conservation planning.
28 *Environmental Management*, **35**, 527-543.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. GIS data used to characterize the planning units in the Guadiana River basin at the catchment scale (the whole upstream drainage area excluding itself) and the subcatchment scale (only the stretch within the planning unit in question). SD denotes Standard Deviation.

Catchment scale	Subcatchment scale
% Land uses ¹ :	% Land uses ¹ :
Urban	Urban
Intensive agriculture	Intensive agriculture
Extensive agriculture	Extensive agriculture
Naturalized	Naturalized
% Geology ² :	% Geology ² :
Actual alluvial deposits	Actual alluvial deposits
Calcareous	Calcareous
Siliceous	Siliceous
Acid volcanic	Acid volcanic
Basic volcanic	Basic volcanic
Mean annual rainfall ³	Mean annual rainfall ³
Precipitation seasonality ³	Precipitation seasonality ³
Mean annual temperature ³	Precipitation in the wettest month ³
Altitude (Average, Maximum and SD) ⁴	Precipitation in the driest month ³
Slope (Average and SD) ⁴	Mean annual temperature ³
FootPrint ⁵	Annual temperature range
Population density	Temperature in the warmest month ³
Drainage area	Temperature in the coldest month ³
	Temperature seasonality ³
	Isothermality ³
	Mean annual evapotranspiration ³
	Altitude (Average, Maximum and SD) ⁴
	Slope (Average and SD) ⁴
	Soil Quality Index ⁶
	Vegetation Quality index ⁶
	Distance to headwater
	Downstream distance to the nearest reservoir
	Distance to Guadiana (main river channel)
	FootPrint ⁵
	Population density ⁶

Data sources:

1 CORINE Land-Cover 1:100.000. Confederación Hidrográfica del Guadiana.

2 Mapa geológico de España 1:1.000.000. Instituto Geológico y Minero de España.

3 WORLDCLIM, Version 1.4. The data is described in Hijmans et al., (2005).

4 SRTM 90 m Digital elevation model from Jarvis et al., (2006).

5. Human footprint. Center for International Earth Science Information Network (CIESIN) at Columbia University (www.ciesin.columbia.edu/wild_areas/)

6. European Environmental Agency. (www.eea.europa.eu).

Table 2. Principal Component Analysis carried out in the environmental data matrix to select the most representative and independent predictors within the study area. The variance explained by each Principal Component (PC) is showed in addition to their respective eigenvalues (in parentheses). The variable with the highest loading within each PC was selected as representative of each of them and used as predictors. * Denotes sub-catchment measured variables in opposition to catchment variables.

	Variance explained	Environmental variable	Factor loading
<i>PC1</i>	24.9 (12.9)	Coldest temperature*	-0.95
<i>PC2</i>	16.2 (8.4)	Average slope*	0.83
<i>PC3</i>	8.5 (4.4)	Average Evapotranspiration*	-0.65
<i>PC4</i>	7.6 (3.9)	Altitude (SD)	-0.79
<i>PC5</i>	6.3 (3.3)	Warmest temperature*	-0.65
<i>PC6</i>	4.2 (2.2)	Area	0.53
<i>PC7</i>	3.5 (1.8)	Siliceous	-0.56
<i>PC8</i>	2.8 (1.4)	Extensive agriculture*	0.41
<i>PC9</i>	2.7 (1.4)	Siliceous*	0.40
Total	76.7 (39.9)		

Table 3. MARS-GLM model performance. The deviance explained indicated the reduction in deviance for each species with respect a null model. The proportion of total deviance accounted for is shown in brackets. The discriminatory power of the model for each species is given through the AUC of the ROC curve (calculated by K-fold re-sampling with its SD in brackets).

Species	Author	Deviance explained	ROC	Prevalence (n=151)
<i>Anaocypris hispanica</i>	Steindachner, 1866	33.6 (0.41)	0.72 (0.11)	0.05
<i>Luciobarbus comizo</i>	Steindachner, 1864	44.7 (0.27)	0.75 (0.15)	0.24
<i>Luciobarbus microcephalus</i>	Almaça, 1967	46.6 (0.25)	0.70 (0.20)	0.31
<i>Luciobarbus sclateri</i>	Günter, 1868	34.6 (0.34)	0.79 (0.21)	0.10
<i>Iberochondrostoma lemmingii</i>	Steindachner, 1866	42.9 (0.30)	0.73 (0.15)	0.28
<i>Pseudochondrostoma willkommii</i>	Steindachner, 1866	53.2 (0.24)	0.70 (0.20)	0.17
<i>Cobitis paludica</i>	Buen, 1930	32.2 (0.02)	0.57 (0.08)	0.66
<i>Salaria fluviatilis</i>	Aso, 1801	47.8 (0.42)	0.69 (0.28)	0.13
<i>Iberocypris alburnoides</i>	Steindachner, 1866	3.8 (0.18)	0.66 (0.12)	0.66
<i>Squalius pyrenaicus</i>	Günter, 1868	25.5 (0.42)	0.69 (0.16)	0.13
Average		36.5 (0.29)	0.71 (0.16)	0.27

1
2
3
4 **Figure 1.** Location and water courses network of the Guadiana River basin. The main
5 river channel is identified with a thicker line.
6


7 **Figure 2.** Decay in upstream connectivity penalties incurred if planning units are not
8 selected.
9

10 **Figure 3.** Trade-off between minimizing boundary length and total area to reserve for
11 different Connectivity Penalties (CP) values. We set the optimal CP (pointed out with
12 an arrow) at a value where a substantial gain in the reduction of reserve's boundary was
13 get for a minimum increase in total area to reserve.
14
15
16
17

18 **Figure 4.** Total area (Mean \pm SE) included in best solutions after 100 runs in Marxan
19 for different target levels. For each target level 10 different Connectivity Penalties (CP)
20 where used (0, 0.001, 0.05, 0.1, 0.3, 0.5, 0.7, 1, 2, and 3) to reduce the potential effect of
21 changes in optimal CPs for each targets on the results.
22
23
24
25

26 **Figure 5.** Effect of Connectivity Penalty (CP) on reserve design. It is shown the best
27 solutions (a-c) and irreplaceability values (d-f) for a target of 10 when setting the CP at
28 three different levels (0, 0.01 and 3). Planning units included in the best solution found
29 after 100 runs in Marxan, are pointer out in grey (a-c). Higher irreplaceability values are
30 drawn in darker colours (d-e). Irreplaceability represents the frequency of selection of
31 each planning unit by running the selection algorithm 100 times. It is only represented
32 the portion of the basin where the best solutions and irreplaceability values appeared.
33
34
35
36
37
38 The figure in the middle corresponds to the trade-off CP value set in Fig. 3.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig.1



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig 2






Fig 3



for Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig 4



py for Review

Fig 5

