


REVIEW


Holocene biogeography of the southwestern European white-toothed shrew (*Crocidura iculisma*, Eulipotyphla) through its fossil record

Ángel C. DOMÍNGUEZ-GARCÍA*  *Aragosaurus-IUCA, Departamento de Ciencias de la Tierra, Facultad de Ciencias, Universidad de Zaragoza, c/ Pedro Cerbuna 12, 50009 Zaragoza, Spain and Departamento de Geodinámica, Estratigrafía y Paleontología, Facultad de Ciencias Geológicas, Universidad Complutense de Madrid, c/ José Antonio Novais, 12, 28040 Madrid, Spain. Email: angelcdo@ucm.es*

Adrián ÁLVAREZ-VENA *Departamento de Geología, Universidad de Oviedo, c/ Jesús Arias de Velasco s/n, 33005 Oviedo, Spain. Email: adrianalvarezvena@geol.uniovi.es*

César LAPLANA *Departamento de Geología, Geografía y Medio Ambiente, Facultad de Ciencias, Universidad de Alcalá, Ctra. A-II km 33.600, 28871 Alcalá de Henares, Spain and Museo Arqueológico y Paleontológico de la Comunidad de Madrid, Plaza de las Bernardas s/n, 28801 Alcalá de Henares, Madrid, Spain. Email: cesar.laplana@uah.es*

Paloma SEVILLA *Departamento de Geodinámica, Estratigrafía y Paleontología, Facultad de Ciencias Geológicas, Universidad Complutense de Madrid, c/ José Antonio Novais, 12, 28040 Madrid, Spain. Email: psevilla@ucm.es*

Jacinto ROMÁN  *Departamento de Biología de la Conservación, Estación Biológica de Doñana – EBD/CSIC, Avda. Américo Vespucio 26, 41092 Sevilla, Spain. Email: jroman@ebd.csic.es*

Josep Francesc BISBAL-CHINESTA *Institut Català de Paleoecologia Humana i Evolució Social, Zona Educacional 4, Campus Sescelades URV, Edifici W3, 43007 Tarragona, Spain and Departament d'Història i Història de l'Art, Universitat Rovira i Virgili, Tarragona, Spain. Email: jbisbal@iphes.cat*

Javier CALZADA *Universidad de Huelva Departamento de Ciencias Integradas, Facultad de Ciencias Experimentales, Universidad de Huelva, Av. Tres de marzo s/n, 21007 Huelva, Spain. Email: javier.calzada@dbasp.uhu.es*

M. Ángeles GALINDO-PELLICENA *Centro Mixto UCM-ISCIII de Evolución y Comportamiento Humanos, C/ Monforte de Lemos 5, 28029 Madrid, Spain. Email: mariangalindo79@gmail.com*

Luis BENÍTEZ DE LUGO ENRICH *Departamento de Prehistoria, Historia Antigua y Arqueología, Facultad de Geografía e Historia, c/ Prof. Aranguren s/n. Edif. Filosofía B, E-28040 Madrid, Spain. Email: luisbeni@ucm.es*

Keywords

Crocidura russula, Iberian Peninsula, interspecific competition, small mammals, Soricidae

*Correspondence

Received: 22 March 2024

Accepted: 8 July 2024

Editor: RG

doi: 10.1111/mam.12374

ABSTRACT

1. *Crocidura iculisma* (= *C. suaveolens*) is a shrew (Soricidae, Eulipotyphla) showing a fragmented distribution limited to the Iberian Peninsula, southern and western France and northwestern Italy. Although it has been in the Iberian Peninsula since the Middle Pleistocene, its biogeographical history remains poorly known.
2. Here, we provide new data on *Crocidura iculisma* from the Castillejo del Bonete site (southern Spain) dated as 3.8–3.6 cal kyr BP. At the same time, we analyse changes in its geographic distribution based on its palaeontological record.
3. We have reviewed thoroughly the Holocene record of *Crocidura iculisma* in southwestern Europe, considering independently the three time intervals according to the stages defined for this geological time series (Greenlandian,

Northgrippian and Meghalayan). Whenever found together in fossil sites, the relative abundance of the two sympatric shrews of the genus *Crocidura* (*Crocidura russula* and *Crocidura iculisma*) was analysed. The data obtained concerning past distribution range and relative abundance were compared with the current status of the species in Iberia to interpret changes through time.

4. The distribution range of the southwestern European white-toothed shrew in the Iberian Peninsula has notably reduced over time until today. The occurrence at Castillejo del Bonete and other palaeontological records located beyond its current distribution range show that it was distributed in several areas of the eastern part and southern interior of Iberia, at least until the second half of the Meghalayan. Relative abundance patterns observed in the fossil assemblages suggest that competitive exclusion by *Crocidura russula* and climatic changes played a crucial role in driving the extirpation of *Crocidura iculisma* from a substantial portion of the Iberian Peninsula.

INTRODUCTION

From a palaeontological standpoint, the study of the Holocene small mammal record in southwestern Europe allows the reconstruction of recent biogeographic processes influenced by climatic evolution, ecological interactions or human impacts (Auffray et al. 1990, Poitevin et al. 1990, Vigne 1992, Cuenca-Bescós et al. 2009, Laplana & Sevilla 2013, Domínguez García et al. 2019, Álvarez-Vena et al. 2023). Particularly for shrews (Soricidae), some research has been done (e.g. Rzebik-Kowalska 1998, Furió 2007, Rofes & Cuenca-Bescós 2011, Moya-Costa & Cuenca-Bescós 2021, Moya-Costa et al. 2023), although few studies have focused on their Holocene palaeobiogeography (Rofes et al. 2018, Domínguez García et al. 2020).

Under the name of the lesser white-toothed shrew, *Crocidura suaveolens* s.l., a species group of tiny shrews, inhabits a wide geographic range in Eurasia, extending continuously from Central Europe to Asia (Palomo et al. 2016). In Western Europe, they are less abundant, particularly in the Iberian Peninsula, where a single species is represented with a small distribution since it is found mainly restricted to areas of Atlantic influence (Rey 2007, Bencatel et al. 2017).

The systematics and taxonomic status of the *Crocidura suaveolens* s.l. group has been considered one of the most problematic within the genus (Hutterer 2005). Mitochondrial DNA studies have found that the white-toothed shrews of Europe belong to two clearly differentiated clades (Dubey et al. 2006, Biedma et al. 2018): one from central Europe, corresponding to the subspecies *Crocidura suaveolens mimula*, and another from the extreme southwest, corresponding to the subspecies *Crocidura suaveolens iculisma*. The latter shows a wide divergence and is placed in a more basal position than other clades and is clearly differentiated from the other

European clade (Dubey et al. 2006, 2007). In parallel, Bannikova et al. (2006), in a study with mitochondrial DNA which did not include individuals from the southwestern European clade, concluded that *Crocidura s. mimula* should be considered a valid species and named *Crocidura mimula*, which is in agreement with the results obtained by Dubey et al. (2006), who suggested that the status of many taxa should be re-evaluated. Recent studies (Wilson & Mittermeier 2018, Burgin et al. 2020) have proposed, as a preliminary solution and in the absence of nDNA analysis, that populations from western and central Europe, Mediterranean islands and southwestern Asia should be classified as *Crocidura gueldenstaedtii*, while *Crocidura suaveolens* should be retained for populations from eastern Europe and Asia. This proposal has also been recently adopted by the IUCN Red List of Threatened Species (Kryštufek & Gazzard 2023). However, the most recent studies, using mitogenomes and nDNA, have concluded that both *Crocidura mimula* and *Crocidura iculisma* are valid species (Gritsyshin et al. 2023, İbiş et al. 2023). Therefore, here we will use *Crocidura iculisma* to refer to the populations of lesser white-toothed shrews of the Iberian Peninsula and Mediterranean France, as this is in agreement with the most recent knowledge. Under this denomination, we have included all previous mentions of *Crocidura suaveolens* or *Crocidura gueldenstaedtii* from the Iberian Peninsula or France.

Fossils assigned to *Crocidura iculisma* have been described in the Iberian Peninsula since the Middle Pleistocene, probably as a descendant of *Crocidura kornfeldi* (Rofes & Cuenca-Bescós 2011), although fossil occurrences are not abundant (Sesé & Sevilla 1996, Arribas 2004, Moya-Costa 2020). During the Pleistocene glacial periods, the southern European peninsulas such as Iberia, Italy and the Balkans operated as refugia for temperate species (Hewitt 1996, Taberlet et al. 1998, Sommer & Nadachowski 2006). Previous works

have recognised a post-glacial diversification and dispersal process of *Crocidura iculisma* from its Iberian refuge to the north fringe of Europe (Biedma et al. 2018, Rofes et al. 2018). Particularly, the Western European populations show a current geographic distribution that is fragmented in various areas due to the evolution of three main lineages in separate glacial sub-refugia during the Quaternary (Biedma et al. 2018). In addition, some authors have also suggested that the use of resource competition with the congeneric species *Crocidura russula* has been a key factor in its biogeographic history, showing the influence of competitive exclusion of *Crocidura iculisma* by *Crocidura russula* in large areas of Europe (Niethammer 1979, Poitevin et al. 1987, Kraft 1998, Biedma et al. 2018, 2020, Bellocq et al. 2023).

The main objective of this study is to reconstruct the Holocene biogeographic history of *Crocidura iculisma*, analyse its changes compared to the present and infer the causes behind these changes. To achieve this goal: 1) we present a new record of *Crocidura iculisma* coming from Castillejo del Bonete site in southeastern Spain; 2) we compiled all known records of the species from Holocene archaeological and palaeontological sites in southwestern Europe; and 3) we conducted a comparative analysis of the relative abundances of *Crocidura iculisma* and *Crocidura russula* between past and modern assemblages.

METHODS

New data on the presence of *Crocidura iculisma*

The shrew remains studied in this work come from Castillejo del Bonete site (Terrinches, Ciudad Real, Spain), one of the most outstanding archaeo-palaeontological sites for studying Holocene small mammals in the Iberian Peninsula. It is a burial mound complex from the Chalcolithic and Bronze Age built on a cave (Benítez de Lugo et al. 2014, 2015, Benítez de Lugo 2018). The sediments of the underlying cave have yielded an exceptionally large and diverse microvertebrate record, providing new valuable data to reconstruct the biogeographical history of some species in Western Europe (Domínguez García et al. 2019, Benítez de Lugo et al. 2020).

Crocidura spp. cranial remains from the cave of Castillejo del Bonete were picked from the fossiliferous sediments excavated during the fieldwork campaign in 2012. The specimens studied here come from the west test pit of Gallery 2 (SWG2, layers UE 26018, UE 26019 and UE 26020), its small mammal assemblage previously described by Domínguez García et al. (2019), who identified a minimum number of 81 individuals (MNI) corresponding to *Crocidura russula*. For a more detailed description of the site and the collecting and sorting methodology for the

recovery of the microvertebrate remains, see Domínguez García et al. (2019) and Benítez de Lugo et al. (2015, 2020).

Concerning the age of this microvertebrate assemblage, the material from SWG2 has been dated as 3,819–3,615 cal years BP (Domínguez García et al. 2019, Benítez de Lugo et al. 2020), thus evidencing that the remains were accumulated during the early Meghalayan (Late Holocene).

Taxonomic identifications were made based on the morphology of diagnostic elements (upper fourth premolar – P4, and second lower molar – m2) following the criteria broadly recognised to discriminate dental remains of *Crocidura iculisma* from those of *Crocidura russula* (Rey & Landin 1973, Saint-Girons et al. 1979, Poitevin et al. 1986, Calzada & Román 2017) and also genetically validated by Biedma et al. (2018). To characterise the size of the *Crocidura* remains here analysed, length and width measurements of the P4 were taken following Reumer (1984). Comparisons were made with specimens from modern localities and Holocene sites from Spain and Mediterranean France. Measurements of the modern referential material were taken on the samples from the collections of the Museum National d'Histoire Naturelle (MNHN, Paris, France) and the Estación Biológica de Doñana (EBD-CSIC, Sevilla, Spain) (Appendix S1).

Review of Holocene record

To reconstruct the biogeographical history of *Crocidura iculisma*, a review of published Holocene records in the Iberian Peninsula and Mediterranean France was conducted. An updated database was built for this purpose, considering the taxonomic and/or chronological precision of each case. Previous publications reviewing the record of Iberian Quaternary fossil mammals (Arribas 2004, López-García 2011) were considered.

All fossil occurrences of *Crocidura iculisma* were compiled in the database and arranged in three intervals according to the stages/subepochs of the Holocene following Walker et al. (2018): Greenlandian/Early Holocene (11.7–8.2 ka), Northgrippian/Middle Holocene (8.2–4.2 ka) and Meghalayan/Late Holocene (4.2 ka till present). The geographic location of these records was compared to the current distribution range of the species (Rey 2007, Palomo et al. 2016, Bencatel et al. 2017) to identify changes that may have during the Holocene in southwestern Europe.

Abundance ratio analysis

We compiled published fossil assemblages in which *Crocidura iculisma* and *Crocidura russula* co-occurred




Fig. 1. Selected fossil specimens from Castillejo del Bonete. *Crocidura iculisma*: left maxilla with A3-M2 in labial (a) and occlusal (a') views (BO12_SWG2_26018_018); *Crocidura russula*: skull with complete dentition in labial (b) and occlusal (b') views (BO18_S2.2G2_229_020), right mandible with complete dentition in posterior (c) and labial (c') views (BO18_S2.2G2_229_014). Arrows indicate the position of P4 in the maxilla. Dashed lines indicate morphological diagnostic features. Scale bars 1 mm.

and the abundance data provided by authors (Asquerino & Woloszyn 1991, Poitevin & S negas 1999, Poitevin et al. 2005, Valenzuela et al. 2009, Guillem Calatayud 2011,  lvarez-Vena et al. 2023, Cernadas-Garrido et al. 2023). Taphonomic information related to the origins of each assemblage was considered, whenever available, to make abundance values with modern assemblages comparable.

In addition, we collected published data on the abundance of the two shrew species from modern populations obtained from owl pellet analyses (Nores 1989, Biedma et al. 2019) and live trapping (Biedma et al. 2020). We also included unpublished data obtained from barn owl pellets recovered at 12 modern localities showing the co-occurrence of both *Crocidura* species. The biogeographical region (Eurosiberian or Mediterranean) in which each site and modern locality are located was also considered.

To test the possible dominance pattern between the two *Crocidura* species, the relative abundance (% MNI) of each was calculated for all the Holocene fossil assemblages in which they were found together, and the resulting values were compared with the relative proportion in which they are represented in modern localities. To check if there are differences between fossil sites and modern localities, a statistical analysis was conducted using the Clopper–Pearson method (Clopper & Pearson 1934) with PAST 4.01 software (Hammer et al. 2001). We calculated multiple-proportion 95% confidence intervals (CI) for the relative abundance of each species in every site and modern population. Following Such eras-Marx et al. (2019), variations were considered statistically significant when CI

values for each species did not overlap between fossil and modern assemblages. Since both *Crocidura* shrews are potentially found in the same habitats and exhibit similar morphology and behaviour, it is reasonable to assume equal catchability by predators (Biedma et al. 2019). Therefore, ecological factors other than selective predation can be argued to explain differences in abundance values between assemblages.

RESULTS AND DISCUSSION

Castillejo del Bonete record

This study adds a new Holocene occurrence of the south-western European white-toothed shrew in Spain, belonging to the west test pit in Gallery 2 (SWG2) of the cave of Castillejo del Bonete site. A maxilla from this assemblage assigned in a previous paper (Dom nguez Garc a et al. 2019) to *Crocidura russula* was found to display the features that are characteristic of *Crocidura iculisma* and have been consequently included in the present study (Fig. 1a).

The morphological traits that allow assigning material to *Crocidura iculisma* according to Rey and Landin (1973), Saint-Girons et al. (1979), Poitevin et al. (1986) and Calzada and Rom n (2017) are the rounded outline in the upper edge of the parastyle on the fourth upper premolar (P4) in labial view, being well developed behind the paracone and with a thick cingulum (Fig. 1a) and the position in occlusal view of its protocone which stands at a distance from the lingual edge of the tooth (Fig. 1a'). On the other hand, in the P4 of *Crocidura russula*, the parastyle

has an angular outline in labial view (Fig. 1b) and the protocone in occlusal view is found at the lingual border (Fig. 1b'); in the lower dentition, the m2 of *Crociodura russula* on labial view displays a distinct constriction in its labial cingulum (Fig. 1c'). Although this constriction is less pronounced in certain remains, no morphotypes of *Crociodura iculisma* showing a straight labial cingulum have been found.

Additionally, the biometrical study supports the taxonomic assignments of the material from Castillejo del Bonete (Fig. 2, Appendix S1). Overall, the P4 in *Crociodura iculisma* is smaller than *Crociodura russula*, except in Punta Lucero III. This is in accordance with the size gradient pattern observed today in the Iberian Peninsula, which reveals that *Crociodura* spp. populations of the Basque Country are considerably larger than in other more southern localities (Rey & Landin 1973, Zabala 1985). The specimen from Castillejo del Bonete assigned to *Crociodura iculisma* clearly falls within the range of variability of this species, being closer to the smaller values obtained in the Cueva del Caballo (6) and Postes Cave (9) samples.

In the same way, the size of *Crociodura russula* from Castillejo del Bonete (1) is closer to the smaller values within the range displayed by the species in the Cueva del Estrecho sample (10).

Biogeography

The review of the Holocene record of *Crociodura iculisma* in the Iberian Peninsula and southern France has provided 21 records, to which the new occurrence from Castillejo del Bonete has now been added. Eleven of these records come from sites that are geographically placed out of the current distribution range of the species, mainly in eastern and southern Spain (Fig. 3, Table 1).

A few of these records may be considered to not be fully reliable, either because the age of the site is imprecise or because no descriptions or figures supporting the taxonomic assignment are given. This is the case of Cueva de Ardales (1), for which a wide temporal range between Late Pleistocene and Late Holocene is assigned to the single remains of *Crociodura iculisma* (Rofes 2023) found in the site. Similarly,




Fig. 2. Bivariate plot comparing the length and width of the P4 (mean ± SD values). 1–2: Castillejo del Bonete; 3–4: Punta Lucero III; 5–6: Cueva del Caballo; 7–8: Cueva de los Mármoles; 9: Postes Cave, 10: Cueva del Estrecho; 11: Mediterranean France; 12, 14: Doñana/Cádiz; and 13: Doñana/Almonte. 1–6: This work (Appendix S1); 7–8: (Asquerino & Woloszyn 1991); 9: (Domínguez-García et al. 2023); and 10–14: this work (Appendix S1).




Fig. 3. Map showing the *Crocidura iculisma* records in southwestern Europe considered in this work. Letters and numbers (ID, Table 1) indicate modern localities and Holocene fossil sites respectively. Fossil occurrences spanning multiple time intervals are denoted by points of various colours following the legend. The current geographic distribution of the species is based on Palomo et al. (2016). Map modified from the GEBCO_2021 Grid (GEBCO Bathymetric Compilation Group 2021).

no radiocarbon date is provided for Cueva de los Mármoles (8), where *Crocidura iculisma* was found in a context linked to the Middle Neolithic based on its archaeological content (Asquerino & Woloszyn 1991). Other sites (2, 3, 4, 7, 12) lack descriptions or figures describing the material (Poitevin et al. 1990, Pemán 1994, Jeannet & Vital 2009) preventing verification of the taxonomic assignment. Apart from these, the remaining records can be considered both taxonomically and chronologically reliable.

The presence of *Crocidura iculisma* in Mesolithic levels of Postes Cave (5), in the Neolithic–Chalcolithic of Cueva de los Mármoles (8), Cueva del Caballo (9) and Portalón (10), and finally, in the Bronze Age of Castillo del Bonete (16) shows its persistence in the southern interior regions of the Iberian Peninsula, at least until the first half of Meghalayan. At the same time, the site of Castro do Zambujal (13) has yielded a record of *Crocidura iculisma* right at the southernmost limit of its current distribution in Portugal. Concerning the records of eastern Spain, most of them belong to sites located in the easternmost mountain ranges of the Iberian and Baetic mountainous systems (6, 11, 15, 19), showing that *Crocidura iculisma* survived there until at least the Iron Age and Roman Period, according to its record in c. 2000 years ago in Alorda Park (18) and Estrets-Racó de Rata (19). For the Eurosiberian region, where the species is still found today, three occurrences have been published,

belonging to the sites of Laminak (2), Punta Lucero III (14) and level 0 of Cueva del Hueso (22).

Relative abundance

A total of nine archaeological and palaeontological sites where *Crocidura iculisma* and *Crocidura russula* co-occurred were compiled, with Castillo del Bonete included in the abundance ratio analysis (Table 2, Fig. 4).

A common pattern emerges from the archaeological and palaeontological assemblages regarding abundance ratios. Whereas *Crocidura russula* showed notably higher abundances than *Crocidura iculisma* in all mentioned sites, at Punta Lucero III level F (Table 2, Fig. 4) this ratio reversed. This indicates that in the areas surrounding the archaeological and palaeontological sites where *Crocidura iculisma* has been found, the species was less common than *Crocidura russula*. This relative abundance pattern coincides with what is observed in current sympatric populations in the Mediterranean and Eurosiberian regions, except for the population studied through live trapping within the Odiel Marshes Natural Reserve (B – Biedma et al. 2020), where *Crocidura iculisma* was found and clearly dominated the tidal marsh environment. The authors of this research interpreted this result as an evolutionary response of *Crocidura iculisma*, indicating its

Table 1. Holocene palaeontological sites and modern localities with *Crocidura iculisma* records in southwestern Europe. The ID refers to Fig. 3. Abbreviations: ES=Spain, FR=France, PT=Portugal

Time interval	ID	Site/Locality	Levels	Age	Reference
Late Pleistocene-Meghalayan/Late Holocene	1	Cueva de Ardales (ES)	Zone 2 – Layers 16–14	29–3.8 cal kyr BP	Rofes (2023)
Late Pleistocene-Greenlandian/Early Holocene	2	Laminak II (ES)	L. I (1–2)	12.5–11.7 cal kyr BP	Pemán (1994)
Greenlandian/Early Holocene	3	Chauve Souris (FR)	BA	±9.9 kyr BP	Jeannet and Vital (2009)
Greenlandian-Northgrippian/Early-Mid Holocene	4	Fontbrégoua (FR)	71–54; 53–31	8.6–8.2 cal kyr BP 6.8–6.2 cal kyr BP	Poitevin et al. (1986, 1990)
	5	Cueva de los Postes (ES)	Stage 3	8.4–7.4 cal kyr BP	Domínguez-García et al. (2023)
	6	Abric de la Falguera (ES)	Phases VIII–VI	8.4–6.3 cal kyr BP	Guillem Calatayud (2006)
	7	L’Abri de la Font-des Pigeons (FR)	L. 19–10	8.6–8.2 cal kyr BP 6.9–6 cal kyr BP	Poitevin et al. (1986, 1990)
Northgrippian/Middle Holocene	8	Cueva de los Mármoles (ES)	Surface, SW, F, SC	–	Asquerino and Woloszyn (1991)
	9	Cueva del Caballo (ES)	UE 16	5.6–5.7 cal kyr BP	Domínguez García (2021)
	10	Portalón (ES)	Funerary – Habitat/Stabling	5.3–4.2 cal kyr BP	Rofes et al. (2021)
	11	Abric de la Falguera (ES)	Phases V–IV	5.1–4.4 cal kyr BP	Guillem Calatayud (2006)
Northgrippian-Meghalayan/ Mid-Late Holocene/	12	Chauve Souris (FR)	15–6C	4.9–2.3 kyr BP	Jeannet and Vital (2009)
Meghalayan/Late Holocene	13	Castro do Zambujal (PT)	–	4.8–3.9 ca kyr BP	Storch and Uerpmann (1976)
	14	Punta Lucero III (ES)	F	4.4–4.1 cal kyr BP	Álvarez-Vena et al. (2023)
	15	Cova de les Tàbegues (ES)	–	4–3 kyr BP	Guillem Calatayud (2002)
	16	Castillejo del Bonete (ES)	UE 26020-26018 SWG2	3.8–3.6 cal kyr BP	This work, Domínguez García et al. (2019)
	17	Lattara (FR)	–	2.4–2.2 BP	Poitevin and Sénégas (1999)
	18	Alorda Park (ES)	–	2.3–2.1 cal kyr BP	Valenzuela et al. (2009), Rofes et al. (2018)
	19	Estrets-Racó de Rata (ES)	2004–2005	2.3–2.1 kyr BP	Guillem Calatayud (2011)
	20	Puits de Lattara (FR)	–	2.1–1.8 kyr BP	Poitevin et al. (2005)
	21	Canet 59 (FR)	–	2 cal kyr BP	Callou and Vigne (2013), Rofes et al. (2018)
Modern	22	Cueva del Hueso (ES)	0	–	Cernadas-Garrido et al. (2023)
	A	Guadalquivir marshes (ES)	–	–	Biedma et al. (2019)
	B	Odiel marshes (ES)	–	–	Biedma et al. (2020)
	C	Cuacos de Yuste (ES)	–	–	This work
	D	Gata (ES)	–	–	This work
	E	Candelario (ES)	–	–	This work
	F	Torres de Aliste (ES)	–	–	This work
	G	Figueruela de Arriba (ES)	–	–	This work
	H	Castro Candelas (ES)	–	–	This work
	I	Forcarei (ES)	–	–	This work
	J	Infiesto (ES)	–	–	Nores (1989)
	K	Avín (ES)	–	–	This work
	L	Partearroyo (ES)	–	–	This work
	M	Carniego (ES)	–	–	This work
	N	Ungo (ES)	–	–	This work
	O	Encima Angulo (ES)	–	–	This work

specialisation to tidal marsh habitats to avoid direct competition with *Crocidura russula*. Relative abundance patterns in the areas where both species co-occur may explain the result of highly asymmetric interspecific competition, with *Crocidura russula* being the dominant species. This dominance may have led to the gradual displacement of

Crocidura iculisma from large areas, pushing it into the most humid and forested habitats, as seen in its current distribution and, in a more extreme way, to the tidal marshes in the Gulf of Cádiz (Niethammer 1979, Poitevin et al. 1987, Román & Ruiz 2003, Biedma et al. 2018, 2019, 2020).

Table 2. List of sites and modern localities included in the abundance relationship analysis. ID referred to Figs 3 and 4. MNI=Minimum number of individuals, *C. i*=*Crocidura iculisma*, *C. r*=*Crocidura russula*

Biogeographic region	ID	Site/locality	MNI <i>C. i</i>	MNI <i>C. r</i>	Total MNI	%MNI <i>C. i</i>	%MNI <i>C. r</i>	Origin of the assemblage	
<i>Palaeontological sites</i>									
Mediterranean	8	Cueva de los Mármoles	29	44	73	39.73	60.27	Indeterminate	
	16	Castillejo del Bonete	1	78	79	1.27	98.73	Generalist carnivore predation	
	17	Lattara	5	13	18	27.78	72.22	Probably barn owl pellets	
	18	Alorda Park	12	352	364	3.3	96.7	Barn owl pellets	
	19	Estrets-Racó de Rata	5	27	32	15.63	84.38	Barn owl pellets	
	20	Puits de Lattara	3	63	66	4.55	95.45	Barn owl pellets	
Eurosiberian	14-F	Punta Lucero III - F	49	9	58	84.48	15.52	Accidental falling and/or nocturnal birds of prey pellets	
	14-C-B	Punta Lucero III - C-B	0	130	130	0	100	Accidental falling and/or nocturnal birds of prey pellets	
	22	Cueva del Hueso	11	60	71	15.49	84.51	Nocturnal birds of prey pellets	
<i>Modern localities</i>									
Mediterranean	A	Guadalquivir marshes	32	341	373	8.58	91.42	Barn owl pellets	
	B*	Odiel marshes	0	17	17	0.00	100	Live trapping	
	A-Tm	Guadalquivir marshes	68	256	324	20.99	79.01	Barn owl pellets	
	B-Tm*	Odiel marshes	Tidal marshes <220m	355	27	382	92.93	7.07	Live trapping
			Tidal marshes						
	C	Cuacos de Yuste	66	2729	2795	2.36	97.64	Barn owl pellets	
	D	Gata	13	510	523	2.49	97.51	Barn owl pellets	
	E	Candelario	38	230	268	14.18	85.82	Barn owl pellets	
	F	Torres de Aliste	5	173	178	2.81	97.19	Barn owl pellets	
	G	Figueruela de Arriba	4	176	180	2.22	97.78	Barn owl pellets	
Eurosiberian	H	Castro Candelas	5	237	242	2.07	97.93	Barn owl pellets	
	I	Forcarei	44	105	149	29.53	70.47	Barn owl pellets	
	J	Infiesto	35	255	290	12.07	87.93	Barn owl pellets	
	K	Avin	9	105	114	7.89	92.11	Barn owl pellets	
	L	Partearroyo	6	174	180	3.33	96.67	Barn owl pellets	
	M	Carniego	6	375	381	1.57	98.43	Barn owl pellets	
	N	Ungo	5	297	302	1.66	98.34	Barn owl pellets	
O	Encima Angulo	2	191	193	1.04	98.96	Barn owl pellets		

The statistical test conducted using the 95% confidence intervals (CI) for the relative abundance of each species does not reveal statistically significant overall differences among fossil sites, nor when compared to modern localities from the Mediterranean and the Eurosiberian regions (Fig. 4). Fossil samples with reduced sizes (*Crocidura* spp. MNI < 35) display wider CIs, overlapping regardless of their biogeographic region between them and with certain modern localities. Excluding these samples, the Cueva de los Mármoles (8) shows a high relative abundance of *Crocidura iculisma*, with CI values indicating significant differences from the rest of the fossil sites, and overlapping only with the sample from Forcarei in Galicia (I), which is under a strong Atlantic influence. This situation could be a consequence of the Middle Holocene age of this assemblage, which is the oldest considered for this analysis. Climate and environment were then distinctly more humid than in later periods in the Mediterranean region of Iberia, under a clear trend of aridification since the Middle Holocene onwards (Jalut et al. 2009; Carrión et al., 2010; Ilvonen et al. 2022).

Consequently, humid habitats would have been more prevalent during this period, favouring the presence of *Crocidura iculisma*. Nevertheless, the precise age and agents involved in the accumulation of this assemblage are unclear, and in need of further chronological and taphonomic analyses to consider this result relevant.

A particularly noteworthy case is that of Punta Lucero III at level F (14-F), as it is the only sample displaying differences among all fossil sites, with CI values only overlapping with those from the Odiel Marshes Natural Reserve obtained by live trapping in tidal marshes (B-Tm*). In the stratigraphic sequence of the site, no *Crocidura iculisma* was found at younger levels C and B, where *Crocidura russula* was represented alone (14-C-B). This has been interpreted as an example of displacement of *Crocidura iculisma* by *Crocidura russula*, linked to the progressively drier and anthropised environments (Álvarez-Vena et al. 2023).

The cases of Castillejo del Bonete (16), Alorda Park (18) and Puits de Lattara (20) are also noteworthy for their reduced CI due to their high sample sizes, absence




Fig. 4. Abundance ratio (%MNI) among *Crocidura iculisma* and *Crocidura russula* in fossil sites and modern localities is listed in Table 2. (*) Indicates live-trapping method. *T. marsh* (tidal marshes) refers to samples obtained from owl roosts located less than 220 m away from the tidal marshes, as well as the sample obtained by live trapping within the tidal marshes. MNI, Minimum number of individuals of *Crocidura* spp. CI, Multiple-proportion 95% confidence intervals for the %MNI obtained using the Clopper–Pearson method (Clopper & Pearson 1934).

of significant differences between them or with most of the modern localities in the Mediterranean and the Eurosiberian regions. The assemblage from Cueva del Hueso (22) displays a similar pattern, showing higher abundance of *Crocidura iculisma* with CI values overlapping with Puits of Lattara but displaying significant differences with Castillejo del Bonete and Alorda Park values. Compared to modern referential data, it overlaps with the relatively closer modern populations in Asturias (J, K) and Galicia (I) under the Atlantic climatic influence, and with the samples from Candelario (E) and the Guadalquivir marshes (A, A-Tm). These results are not influenced by differences in sample size nor by their taphonomic origin, given that the accumulation of microvertebrate remains is known to be the result of the accumulation of barn owl pellets, except at Castillejo del Bonete (Table 2). Even though the latter assemblage comes from the activity of generalist carnivore predators (Domínguez García et al. 2019), it has had no influence on the abundance relationship between these two shrew species.

Holocene retraction of the geographic range of *Crocidura iculisma*

Our review shows that the southwestern European white-toothed shrew underwent a significant reduction in its

geographic range during the Holocene. Although fossil occurrences of *Crocidura iculisma* are relatively uncommon, they are enough to show that it still had several populations in the Mediterranean biogeographical region of Spain, even in the Late Holocene. In Mediterranean France, the species has records dating since the Early Holocene and is still extant in the region (Fig. 3). According to our analysis, this biogeographic process could be explained by the combined action of the competitive exclusion with the greater white-toothed shrew and habitat change linked to climatic evolution.

In this sense, since *Crocidura iculisma* has been able to withstand the competitive pressure of *Crocidura russula* in the Mediterranean biogeographic region only in relatively wetter habitats (Poitevin et al. 1987, Román & Ruiz 2003, Biedma et al. 2018, 2019, 2020), it seems reasonable to infer more humid climatic conditions than at present as the conditions in which the Mediterranean fossil assemblages included in this study were formed.

The current Mediterranean climate is characterised by rainy winters and dry summers; the differences between these two seasons being more pronounced in the southern areas and eastern coasts of Iberia (Fick & Hijmans 2017). Throughout the Holocene, the climate in the western Mediterranean realm has been modulated by a series of long-term trends with superimposed short and less intense

drought events (Sánchez Goñi et al. 2008, Jalut et al. 2009, Cacho et al. 2010, Pérez-Obiol et al. 2011, Schirrmacher et al. 2020, Ilvonen et al. 2022). These studies show overall, the existence during the Early and Middle Holocene of an initial relatively humid phase, between 8000 and 6000 years, during which the highest Holocene values of precipitation were reached, known as the Holocene climatic optimum. Posteriorly, after a transitional phase, the increasing aridity trend of the Late Holocene initiated extending until today, being particularly perceptible in the southern regions of Iberia. This provides the scenario in which the wetter conditions of the first half of the Holocene allowed the survival of some *Crocidura iculisma* populations in the eastern and interior areas of the Iberian Peninsula, which persisted at least until the Iron Age–Roman Period transition (Alorda Park and Estrets-Racó de Rata).

Focusing on the statistically significant results of %MNI of *Crocidura* spp., excluding samples influenced by reduced sizes or uncertainties associated with the age of the assemblages mentioned above, *Crocidura iculisma* represents in the fossil samples from the Mediterranean region between 1.27 and 4.55 %MNI (16, 18, 20), similar to those from most modern samples (Table 2, Fig. 4). This suggests that the species was under competitive pressure from *Crocidura russula* in the area around the sites already in the Bronze Age and Iron Age – Roman Period in the Meghalayan. At the same time in Mediterranean France, the southwestern European white-toothed shrew has survived until today, probably because of milder Mediterranean climate and the availability of suitable habitats, while in southern and eastern Iberia, the increasing aridification of climate may have caused the final extirpation of the species, since it is less adapted to face warmer and drier conditions.

Previous studies have already hypothesised that past interspecific competition between the two *Crocidura* species determined their current geographic distribution and recent population dynamics (Poitevin et al. 1987, Biedma et al. 2018, 2019, 2020). Here, we provide for the first time data supporting that this ecological process could have been at work since at least the Meghalayan/Late Holocene in the populations of these two species occupying southwestern Europe.

In addition to the review conducted for this paper, several studies show that *Crocidura russula* is commonly the only species of the genus represented in Holocene Iberian sites. This holds true for areas with a temperate climate (Pemán 1990, Cuenca-Bescós et al. 2009, López-García et al. 2011, Rofes et al. 2013) as well as a Mediterranean climate (e.g. Cardoso et al. 1996, Guillem-Calatayud 1999, 2010, López-García et al. 2010, Bañuls-Cardona et al. 2017, Pimenta et al. 2017, Daura et al. 2019,

Domínguez García et al. 2020) and can be related to the dominance pattern that may have caused the extirpation from most of Iberian territory of *Crocidura iculisma* from up to the present day. However, since *Crocidura iculisma* is much less abundant than *Crocidura russula* both in fossil sites and in the present, and as both species are difficult to distinguish using their fragmentary dental remains, this work exposes the consideration that the western European white-toothed shrew may have often gone unnoticed in fossil assemblages dominated by *Crocidura russula*, generating significant uncertainty about its past distribution. An example of this is given by the maxilla from Castillejo del Bonete included in this work, which was misidentified in a previous study of its small mammal assemblage (Domínguez García et al. 2019).

Some molecular clock studies have estimated that *Crocidura russula* colonised the Iberian Peninsula from Africa during the Late Pleistocene 11–80 kyr (Brändli et al. 2005, Cosson et al. 2005), although an earlier date of arrival is proposed by Biedma et al. (2018), who place this event at the end of Middle Pleistocene (41–200 kyr ~ 126 kyr). When palaeontological data are considered, evidence of an earlier date for its arrival is available, since the oldest Iberian occurrences suggest a minimum age of 300–350 kyr for its presence in Iberia (López Martínez 1980, Sesé & Sevilla 1996, Sesé et al. 2011, 2016, Laplana et al. 2015). However, most of these first records are taxonomically uncertain, since they are based on few fragmentary remains assigned either to *Crocidura* aff. *russula* or *Crocidura* cf. *russula*, and until they are checked there is no solid evidence of the presence of *Crocidura russula* as early as the Middle Pleistocene. On the other hand, several sites from the Middle–Late Pleistocene transition, MIS 5 (Marine Isotope Stage 5), dated 120–80 kyr have yielded more reliable records of *Crocidura russula*, showing the species was widely distributed in the Mediterranean biogeographic region of Iberia (Laplana et al. 2013, López-García et al. 2016, 2022). This latter scenario is more consistent with genetic divergence results and within the maximum age given by Brändli et al. (2005) and Cosson et al. (2005) and the mean date proposed by Biedma et al. (2018).

Despite the new record from Castillejo del Bonete and its importance in the analysis of the biogeography of *Crocidura iculisma* here exposed, our study leaves open some questions. Although our results support the existence of an interspecific competitive relationship between both *Crocidura* species during the Late Holocene, the moment in which it started and how this ecological process might have taken place in the Pleistocene needs to be analysed in more detail. In addition, the morphological and genetic relationships between the modern populations and the fossil specimens are still unknown. The analysis of ancient

DNA (aDNA) in Quaternary small mammals offers valuable information to improve our knowledge of taxonomy, phylogenetic relationships, palaeobiogeographic and palaeoecological studies (Woods et al. 2017, Baca et al. 2020, 2023, Alfaro-Ibáñez et al. 2023, Domínguez-García et al. 2024). This approach could also contribute significantly to advancing this line of research. Indeed, the results presented here leave open further avenues of research, such as extending the temporal range to the Late Pleistocene *Crocidura* records and, including other approaches such as morphometric and molecular analyses, to gather more evidence that might be relevant to clarify the biogeographic history of the *Crocidura iculisma*/*Crocidura russula* relationship in southwestern Europe.

Finally, given that *Crocidura iculisma* is a rare species showing a highly fragmented distribution range, we find it pertinent to discuss some implications for its conservation status. According to the IUCN Red List of Threatened Species, it is classified as least concern (LC) (*Crocidura gueldenstaedtii* in Kryštufek & Gazzard 2023) due to its wide Eurasian range. However, it is classified as near threatened (NT) in the French Red List (MNHN & OFB, 2003–2023), endangered (EN) in Portugal (Calzada et al. 2023) and data deficient (DD) in Spain (Palomo et al. 2007). In the context of global warming, our results reinforce the idea that the *Crocidura iculisma* populations from the Mediterranean climate areas are vulnerable due to habitat and climate change. Its local potential of extinction is enhanced by competition with *Crocidura russula*, which is better adapted to drier and warmer conditions. Conservation actions concerning the habitat where this species is known to be present should be implemented.

As a final remark, this study shows the relevance of palaeontological data as a continuous source of new valuable data to assess the conservation status and threats influencing the species distribution and abundance. Hopefully, it will encourage researchers to carry out further studies focused on *Crocidura* spp. in other regions of southwestern Europe including its fossil record as a valuable source of information.

CONCLUSIONS

The biogeographic history of *Crocidura iculisma* in southwestern Europe is marked by a significant reduction in its distribution range in the Iberian Peninsula during the Holocene.

Our results support the possibility of a relevant role played by interspecific competition and displacement by *Crocidura russula* in explaining its biogeographic dynamics and habitat specialisation. Climatic factors, characterised by increasing aridity over the last 4,000 years, seem to have been decisive in this process so that the reduction

or the disappearance of the humid habitats from the southern and eastern regions of Iberia that once allowed it to withstand competitive pressure with *Crocidura russula* in Mediterranean environments may explain its current low abundance and patchy distribution.

The extent and precision with which palaeontological data contribute to understanding long-term mammal species dynamics are constrained by the nature of these data. Therefore, the similarity in the morphology of both *Crocidura* spp. has put into evidence taxonomic issues regarding the differentiation between both species in fossil small mammal assemblages, restricting the reconstruction of relative abundance ratios and limiting the extent to which their long-term biogeographical history can be reconstructed. Further research complemented with modern techniques will surely help to overcome these issues and achieve a more precise history of these species in the Quaternary of southwestern Europe.

ACKNOWLEDGEMENTS

We would like to thank the following curators for granting access to modern shrew material under their care: C. Urdiales Alonso (EBD-CSIC) and Violaine Nicolas-Colin (MNHN).

FUNDING

ACDG is funded by a postdoctoral Margarita Salas contract (CT18/22) funded by the European Union 'NextGenerationEU/PRTR' and received support from the European SYNTHESYS+ Transnational Access grant (FR-TAF_Call4_007) and the Complutense University of Madrid and Banco Santander by a Research Stay Grant (UCM2020-EB25/20). This is a contribution of the Research Groups UNIZAR E18-23R Aragosaurus and UCM 970827 on Quaternary Ecosystems. This work was partially supported by projects SBPLY/21/180801/000031, SBPLY/22/180801/000042 and SBPLY/23/180801/000006 funded by the Junta de Comunidades Castilla La Mancha, Terrinches City Council and E2IN2.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflict of interest to disclose which could have appeared to influence the work reported in this paper.

DATA AVAILABILITY STATEMENT

Data used for the research described in the article are provided in the main text and Appendix S1 submitted with the paper.

REFERENCES

- Alfaro-Ibáñez MP, Cuenca-Bescós G, Bover P, González Morales M, Straus LG (2023) Implications of population changes among the Arvicolinae (Rodentia, Mammalia) in El Mirón Cave (Cantabria, Spain) for the climate of the last c. 50,000 years. *Quaternary Science Reviews* 315: 108234.
- Álvarez-Vena A, Marín-Arroyo AB, Álvarez-Lao DJ, Laplana C, Arriolabengoa M, Ballesteros D, Aranburu A, Bilbao P, Astorqui Á, Díaz-Casado Y (2023) Mammalian turnover as an indicator of climatic and anthropogenic landscape modification: a new Meghalayan record (Late Holocene) in northern Iberia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 616: 111476.
- Arribas O (2004) *Fauna y Paisaje de los Pirineos en la era glaciár*. Lynx Edicions, Bellaterra, Barcelona, Spain.
- Asquerino MD, Woloszyn BW (1991) Insectívoros y quirópteros del yacimiento neolítico “Cueva de los Mármoles” (Prov. de Córdoba, España). *Estudios de Prehistoria Cordobesa* 5: 43–68.
- Auffray J-C, Vanlerberghe F, Davidian JB (1990) The house mouse progression in Eurasia: a palaeontological and archaeozoological approach. *Biological Journal of the Linnean Society* 41: 13–25.
- Baca M, Popović D, Baca K, Lemanik A, Doan K, Horáček I et al. (2020) Diverse responses of common vole (*Microtus arvalis*) populations to Late Glacial and Early Holocene climate changes – evidence from ancient DNA. *Quaternary Science Reviews* 233: 106239.
- Baca M, Popović D, Agadzhanian AK, Baca K, Conard NJ, Fewlass H et al. (2023) Ancient DNA of narrow-headed vole reveal common features of the Late Pleistocene population dynamics in cold-adapted small mammals. *Proceedings of the Royal Society B: Biological Sciences* 290: 20222238.
- Bannikova AA, Lebedev VS, Kramerov DA, Zaitsev MV (2006) Phylogeny and systematics of the *Crocidura suaveolens* species group: corroboration and controversy between nuclear and mitochondrial DNA markers/ Phylogénie et systématique du groupe d'espèces *Crocidura suaveolens*: coordination et contradiction des marqueurs nucléaire et mitochondriaux de l'ADN. *Mammalia* 70(1–2): 106–119.
- Bañuls-Cardona S, López-García JM, Morales Hidalgo JI, Cuenca-Bescós G, Vergès JM (2017) Lateglacial to Late Holocene palaeoclimatic and palaeoenvironmental reconstruction of El Mirador cave (Sierra de Atapuerca, Burgos, Spain) using the small-mammal assemblages. *Palaeogeography, Palaeoclimatology, Palaeoecology* 471: 71–81.
- Belloq JGD, Fornůšková A, Ďureje L, Bartáková V, Daniszová K, Dianat M et al. (2023) First record of the greater white-toothed shrew, *Crocidura russula*, in the Czech Republic. *Journal of Vertebrate Biology* 72: 23047.
- Bencatel J, Álvares F, Moura AE, Barbosa AM (2017) *Atlas de Mamíferos de Portugal*. Universidade de Évora, Évora, Portugal.
- Benítez de Lugo L (2018) Rituales funerarios neolíticos, calcolíticos y de la Edad del Bronce en la provincia de Ciudad Real: Cerro Ortega (Villanueva de la Fuente) y Castillejo del Bonete (Terrinches). *Anejos a Cuadernos de Prehistoria y Arqueología* 3: 153–168. Madrid.
- Benítez de Lugo L, García HJÁ, Martín SF, Trujillo EM, Sierra JM, Zumajo NP, Lloret CO, Rodríguez AM, Salazar-García DC (2014) Castillejo del Bonete (Terrinches, Ciudad Real): un complejo tumular prehistórico de la cultura de las Motillas en el Alto Guadalquivir. *Menga: Revista de Prehistoria de Andalucía* 5: 151–173.
- Benítez de Lugo L, Zumajo NP, García HJÁ, Bermejo RMB, Sánchez MB, Blain H-A et al. (2015) Paleocología y cultura material en el complejo tumular prehistórico de Castillejo del Bonete (Terrinches, Ciudad Real). *Menga: Revista de Prehistoria de Andalucía* 6: 113–140.
- Benítez de Lugo L, Galindo-Pellicena MÁ, Domínguez García AC, Laplana Conesa C, Herreros GM, Zumajo NP, Romero AP, Sánchez JLF, Salazar-García DC (2020) Fauna e industria sobre materia dura de origen animal del lugar sagrado de la Cultura de las Motillas: Castillejo del Bonete (Terrinches, Ciudad Real). *Cuadernos de Prehistoria y Arqueología de la Universidad Autónoma de Madrid* 46: 45–77.
- Biedma L, Román J, Calzada J, Friis G, Godoy JA (2018) Phylogeography of *Crocidura suaveolens* (Mammalia: Soricidae) in Iberia has been shaped by competitive exclusion by *C. russula*. *Biological Journal of the Linnean Society* 123: 81–95.
- Biedma L, Román J, Godoy JA, Calzada J (2019) Using owl pellets to infer habitat associations and clarify the regional distribution of a cryptic shrew. *Journal of Zoology* 308: 139–148.
- Biedma L, Calzada J, Godoy JA, Román J (2020) Local habitat specialization as an evolutionary response to interspecific competition between two sympatric shrews. *Journal of Mammalogy* 101: 80–91.
- Brändli L, Handley L-JL, Vogel P, Perrin N (2005) Evolutionary history of the greater white-toothed shrew (*Crocidura russula*) inferred from analysis of mtDNA, Y, and X chromosome markers. *Molecular Phylogenetics and Evolution* 37: 832–844.
- Burgin CJ, Wilson DE, Mittermeier RA, Rylands AB, Lacher TE, Sechrest W (2020) *Illustrated Checklist of the Mammals of the World. Volume 2: Eulipotyphla to Carnivora*. Lynx Edicions, Barcelona, Spain.
- Cacho I, Valero-Garcés BL, González-Sampériz P (2010) Revisión de las reconstrucciones paleoclimáticas en la Península Ibérica desde el último periodo glacial. In: Pérez FF, Boscolo R (eds) *Clima en España: Pasado*

- presente y futuro. Informe de evaluación del cambio climático regional, CLIVAR. Ministerio de Medio Ambiente y Medio Rural y Marino: Ministerio de Ciencia e Innovación (MICINN), Spain.
- Callou C, Vigne JD (2013) *Analyse de la faune de deux silos gallo-romains a Canet-en-Roussillon. Vestiges d'un habitat de l'époque romaine et d'un lieu funéraire du haut Moyen Age*. Pyrénées-Orientales (66) Canet-en-Roussillon, Puig del Baja I (fouilles de 1992, 1994 et 2001), 76–81. INRAP, Rapport final d'opération.
- Calzada J, Román J (2017) *EgaEduca. Una herramienta didáctica de investigación con egagrópilas*.
- Calzada J, Mathias ML, Paupério J, Román J (2023) *Crocidura suaveolens* musarinho- de-dentes-brancos. In: Mathias ML, Fonseca C, Rodrigues L, Grilo C, Lopes-Fernandes M, Palmeirim JM, Santos-Reis M et al. (eds) *Livro Vermelho dos Mamíferos de Portugal Continental. FCiências.ID*. ICNF, Lisboa, Portugal.
- Cardoso JL, Antunes MT, Mein PR (1996) Pequenos mamíferos do povoado pré-histórico de Leceia (Oeiras). *Estudos Arqueológicos de Oeiras* 6: 121–133.
- Carrión JS, Fernández S, González-Sampériz P, Gil-Romera G, Badal E, Carrión-Marco Y, López-Merino L, López-Sáez JA, Fierro E, Burjachs F (2010) Expected trends and surprises in the Lateglacial and Holocene vegetation history of the Iberian Peninsula and Balearic Islands. *Review of Palaeobotany and Palynology* 162(3): 458–475.
- Cernadas-Garrido A, Álvarez-Vena A, Álvarez-Lao D (2023) Palaeobiological study of the Holocene micromammal assemblage from the Cueva del Hueso (Castrillón, Asturias, NW Spain). *Spanish Journal of Palaeontology* 38: 235–254.
- Clopper CJ, Pearson ES (1934) The use of confidence or fiducial limits illustrated in the case of the binomial. *Biometrika* 26: 404–413.
- Cosson J-F, Hutterer R, Libois R, Sarà M, Taberlet P, Vogel P (2005) Phylogeographical footprints of the Strait of Gibraltar and Quaternary climatic fluctuations in the western Mediterranean: a case study with the greater white-toothed shrew, *Crocidura russula* (Mammalia: Soricidae). *Molecular Ecology* 14: 1151–1162.
- Cuenca-Bescós G, Straus LG, Morales MRG, Pimienta JCG (2009) The reconstruction of past environments through small mammals: from the Mousterian to the Bronze Age in El Mirón Cave (Cantabria, Spain). *Journal of Archaeological Science* 36: 947–955.
- Daura J, Sanz M, Oms FX, Pedro M, Martínez P, Mendiola S et al. (2019) Deciphering Neolithic activities from a Cardial burial site (Cova Bonica) on the western Mediterranean coast. *Journal of Archaeological Science: Reports* 23: 324–347.
- Domínguez García ÁC (2021) Paleobiogeografía y evolución de las asociaciones de micromamíferos en la península ibérica durante el Holoceno Influencias climáticas y antrópicas.
- Domínguez García ÁC, Laplana C, Sevilla P, Blain H-A, Zumajo NP, Benitez de Lugo Enrich L (2019) New data on the introduction and dispersal process of small mammals in southwestern Europe during the Holocene: Castillejo del Bonete site (southeastern Spain). *Quaternary Science Reviews* 225: 106008.
- Domínguez García ÁC, Laplana C, Sevilla P (2020) Early reliable evidence of the Etruscan shrew (*Suncus etruscus*) in southwestern Europe during ancient times. Reconstructing its dispersal process along the Mediterranean Basin. *Quaternary Science Reviews* 250: 106690.
- Domínguez-García ÁC, Laplana C, Sevilla P, Álvarez-Vena A, Collado Giraldo H (2023) Small mammals of the Holocene sequence of Postes Cave (SW Spain): biogeographic and palaeoenvironmental implications for southwestern Iberia. *Historical Biology* 35: 483–497.
- Domínguez-García ÁC, Utge J, Larrue C, Moclán A, Kibiri Alaoui M et al. (2024) First asserted record of the house mouse in Morocco: application of a multidisciplinary approach to the site of Rirha (5th–1st c. BC). *Archaeol. Anthropological Science* 16: 93.
- Dubey S, Zaitsev M, Cosson J-F, Abdulkadier A, Vogel P (2006) Pliocene and Pleistocene diversification and multiple refugia in a Eurasian shrew (*Crocidura suaveolens* group). *Molecular Phylogenetics and Evolution* 38: 635–647.
- Dubey S, Cosson J, Magnanou E, Vohralík V, Benda P, Frynta D, Hutterer R, Vogel V, Vogel P (2007) Mediterranean populations of the lesser white-toothed shrew (*Crocidura suaveolens* group): an unexpected puzzle of Pleistocene survivors and prehistoric introductions. *Molecular Ecology* 16(16): 3438–3452.
- Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* 37: 4302–4315.
- Furió M (2007) Los Insectívoros (Soricomorpha, Erinaceomorpha, Mammalia) del Neógeno Superior del Levante Ibérico.
- GEBCO Bathymetric Compilation Group (2021) *The GEBCO_2021 Grid – a Continuous Terrain Model of the Global Oceans and Land*. NERC EDS British Oceanographic Data Centre NOC, Southampton, UK.
- Gritsyshin VA, Lisenkova AA, Speranskaya AS, Artyushin IV, Sheftel BI, Lebedev VS, Bannikova AA (2023) Multilocus analysis of phylogenetic relationships in the *Crocidura suaveolens* Ssensu Lato species complex: a comparison with mitochondrial data. *Doklady Biological Sciences* 509: 128–134.
- Guillem Calatayud PM (2002) Aproximación al marco geográfico de la Valltorta y algunas consideraciones sobre la evolución del paisaje Holoceno. In: Martínez Valle R, Villaverde V (eds) *La Cova dels Cavalls en el Barranc de la Valltorta, Monografias del Instituto de Arte Rupestre*, 33–47. Museu de la Valltorta (Tirig), Valltorta, Spain.

- Guillem Calatayud PM (2006) Los micromamíferos (Insectivora, Rodentia y Chiroptera) del Abric de la Falguera, un pequeño fragmento de la historia de la vida. In: García Puchol O, Aura Tortosa JE (eds) *El abric de la Falguera (Alcoi, Alacant): 8.000 años de ocupación humana en la cabecera del río de Alcoi*, 158–167. Museo Arqueológico de Alicante-MARQ, Alicante, Spain.
- Guillem Calatayud PM (2010) Los micromamíferos (rodentia, insectivora y chiroptera) del Cingle del Mas Cremat. El Cingle del Mas Cremat (Portell de Morella, Castellón). Un asentamiento en altura con ocupaciones del Mesolítico Reciente al Neolítico Final: 126–145.
- Guillem Calatayud P (2011) Els paisatges ramaders en època ibèrica. Una reconstrucció a partir dels micromamífers. In: Valenzuela-Lamas S, Padrós N, Belarte MC, Sanmartí J (eds) *Economia agropecuària i canvi social a partir de les restes bioarqueològiques. El primer mil·lenni aC a la Mediterrània occidental*, 117–121. Universitat de Barcelona, Calafell, Spain.
- Guillem-Calatayud PM (1999) Los micromamíferos (Rodentia, Insectivora y Chiroptera) de la secuencia Holocena de la Cova de les Cendres y Cova Bolomini. *Saguntum-PLAV Extra-2*: 31–36.
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Paleontologia Electronica* 4: 9.
- Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society* 58: 247–276.
- Hutterer R (2005) Order Soricomorpha. In: Wilson DE, Reeder DM (eds) *Mammal Species of the World: A Taxonomic and Geographic Reference*, 220–311. Johns Hopkins University Press, Baltimore, MD, USA.
- İbiş O, Koepfli K-P, Özcan S, Tez C (2023) Whole mitogenomes of Turkish white-toothed shrews, genus *Crocidura* (Eulipotyphla: Soricidae), with new insights into the phylogenetic positions of *Crocidura leucodon* and the *Crocidura suaveolens* group. *Organisms Diversity & Evolution* 23: 221–241.
- Iivonen L, López-Sáez JA, Holmström L, Alba-Sánchez F, Pérez-Díaz S, Carrión JS et al. (2022) Spatial and temporal patterns of Holocene precipitation change in the Iberian Peninsula. *Boreas* 51: 776–792.
- Jalut G, Dedoubat JJ, Fontugne M, Otto T (2009) Holocene circum-Mediterranean vegetation changes: climate forcing and human impact. *Quaternary International* 200: 4–18.
- Jeannot M, Vital J (2009) La grotte de la Chauve-Souris à Donzère (Drôme, France): L'environnement holocène par la méthode quantitative appliquée aux microvertébrés. *Essai de définition et application. Revue de Paléobiologie* 28: 139.
- Kraft R (1998) Ehemalige und aktuelle Verbreitung von Hausspitzmaus, *Crocidurus russula* (Hermann, 1780), und Gartenspitzmaus, *Crocidura suaveolens* (Pallas, 1811), in Bayern. In: *Bonner Zoologische Beiträge: Herausgeber, Vol. 49*, 115–129. Zoologisches Forschungsinstitut Und Museum Alexander Koenig, Bonn, Germany.
- Kryštufek B, Gazzard A (2023) IUCN Red List of Threatened Species: *Crocidura gueldenstaedtii*. e.T29653A221737608. <https://doi.org/10.2305/IUCN.UK.2023-1.RLTS.T29653A221737608.en>
- Laplana C, Sevilla P (2013) Documenting the biogeographic history of *Microtus cabreræ* through its fossil record. *Mammal Review* 43: 309–322.
- Laplana C, Blain H-A, Sevilla P, Arsuaga JL, Baquedano E, Pérez-González A (2013) Un assemblage de petits vertébrés hautement diversifié de la fin du MIS 5 dans un environnement montagnard au Centre de l'Espagne (Cueva del Camino, Pinilla del Valle, communauté autonome de Madrid). *Quaternaire. Revue de l'Association française pour l'étude du Quaternaire* 24: 207–216.
- Laplana C, Herráez E, Yravedra J, Báez S, Rubio-Jara S, Panera J, Rus I, Pérez-González A (2015) Biocronología de la Terraza Compleja de Butarque del río Manzanares en el Estanque de Tormentas al sur de Madrid (España). *Estudios Geológicos* 71: e028.
- López Martínez N (1980) Los micromamíferos (Rodentia, Insectivora, Lagomorpha y Chiroptera), del sitio de ocupación achelense de Áridos-1 (Arganda, Madrid). In: Santonja M, López Martínez N, Pérez González A (eds) *Ocupaciones achelenses en el Valle del Jarama*, 161–202 Excma. Diputación Provincial de Madrid, Madrid, Spain.
- López-García JM (2011) *Los micromamíferos del Pleistoceno superior de la Península Ibérica: Evolución de la diversidad taxonómica y cambios paleoambientales y paleoclimáticos*. Editorial Académica Española, Saarbrücken.
- López-García JM, Blain H-A, Allué E, Bañals S, Bargalló A, Martín P et al. (2010) First fossil evidence of an “interglacial refugium” in the Pyrenean region. *Naturwissenschaften* 97: 753–761.
- López-García JM, Blain H-A, Cuenca-Bescós G, Alonso C, Alonso S, Vaquero M (2011) Small vertebrates (Amphibia, Squamata, Mammalia) from the late Pleistocene-Holocene of the Valdavara-1 cave (Galicia, northwestern Spain). *Geobios* 44: 253–269.
- López-García JM, Fernández-García M, Blain H-A, Sanz M, Daura J (2016) MIS 5 environmental and climatic reconstruction in northeastern Iberia using the small-vertebrate assemblage from the terrestrial sequence of Cova del Rinoceront (Castelldefels, Barcelona). *Palaeogeography, Palaeoclimatology, Palaeoecology* 451: 13–22.
- López-García JM, Fagoaga A, Nabais M, Póvoas L, Zilhão J (2022) Late Quaternary (MIS 5a-5b) climate and environments of western Iberia inferred from the small-mammal assemblage of Gruta da Oliveira, Torres Novas, Portugal. *Palaeogeography, Palaeoclimatology, Palaeoecology* 603: 111194.

- MNHN & OFB (ed) (2003–2023) *Sheet of Crocidura gueldenstaedtii* (Pallas, 1811). Inventaire National du Patrimoine Naturel (INPN), Paris, France. https://inpn.mnhn.fr/espece/cd_nom/899768[Ed].
- Moya-Costa R (2020) *Estudio multidisciplinar y reconstrucción de los sorícidos (Eulipotyphla, 1484 Mammalia) del Cuaternario de la Península Ibérica: sistemática, biometría, 1485 paleoambientes, esmalte dental y 3D*. PhD dissertation. Universidad Zaragoza.
- Moya-Costa R, Cuenca-Bescós G (2021) The Quaternary Soricids (Eulipotyphla, Mammalia) fossil record of the Iberian Peninsula. *Ciências da Terra Procedia* 1: 80–83.
- Moya-Costa R, Cuenca-Bescós G, Rofes J (2023) The shrews (Soricidae, Mammalia) of the Early and Middle Pleistocene of Gran Dolina (Atapuerca, Spain): reassessing their paleontological record in the Iberian Peninsula. *Quaternary Science Reviews* 309: 108093.
- Niethammer J (1979) Arealveränderungen bei arten der spitzmausgattung *Crocidura* in der bundesrepublik Deutschland.
- Nores C (1989) Variación temporal y espacial de micromamíferos: determinación mediante análisis de egagrópilas de *Tyto alba*.
- Palomo LJ, Gisbert J, Blanco JC (eds) (2007) *Atlas y libro rojo de los mamíferos terrestres de España*. Dirección General para la Biodiversidad –SECEM-SECEMU, Madrid.
- Palomo LJ, Kryštufek B, Amori G, Hutterer R (2016) *Crocidura suaveolens*. eIUCN Red List of Threatened Species: e.T29656A22296429.
- Pemán E (1990) *Los micromamíferos de la cueva de Amalda y su significación. Comentarios sobre Pliomys lenki (Heller, 1930) (Rodentia, Mammalia)*, 225–238. Eusko Ikaskuntza, Donostia-San Sebastian, Spain.
- Pemán E (1994) Los micromamíferos de Laminak 11 (Berriatua, Bizkaia). *Kobie* 21: 225–233.
- Pérez-Obiol R, Jalut G, Julià R, Pélachs A, Iriarte MJ, Otto T, Hernández-Beloqui B (2011) Mid-Holocene vegetation and climatic history of the Iberian Peninsula. *The Holocene* 21: 75–93.
- Pimenta C, Monteiro M, Miranda M, Sousa A (2017) Small vertebrates from Chalcolithic contexts in Penedo do Lexim (Mafra, Lisboa): the case study of “Abrigo Locus 3.”
- Poitevin F, Sénégas F (1999) Les micromammifères du site de Lattara. In: Py M (ed) *Lattara Recherche sur le quatrième siècle avant notre ère à Lattes. Mélanges d'histoire et d'archéologie de Lattes*, 635–638. CNRS, Lattes, Paris.
- Poitevin F, Catalan J, Fons R, Croset H (1986) Biologie évolutive des populations Ouest-Européennes de Crocidures. I: Critères d'identification et répartition biogéographique de *Crocidura russula* (Hermann, 1780) et *Crocidura suaveolens* (Pallas 1811). *Revue d'Écologie (la Terre et la Vie)* 41: 299–314.
- Poitevin F, Catalan J, Fons R, Croset H (1987) Biologie évolutive des populations Ouest-Européennes de Crocidures (Mammalia, Insectivora). II. — Ecologie comparée de *Crocidura russula* Hermann, 1780 et de *Crocidura suaveolens* Pallas, 1811 dans le midi de la France et en Corse: rôle probable de la compétition dans le partage des milieux. *Revue d'Écologie (La Terre et La Vie)* 42: 39–58.
- Poitevin F, Bayle P, Courtin J (1990) Mise en place des faunes de micromammifères (rongeurs, insectivores) dans la région méditerranéenne française au Post-glaciaire. *Vie Milieu* 40: 144–149.
- Poitevin F, Martínez Garcia E, Cassaing J, Senegas F (2005) Les micromammifères des puits de Lattara. *Lattara* 18: 307–312.
- Reumer JWF (1984) Ruscinian and early Pleistocene Soricidae (Insectivora, Mammalia) from Tegelen (the Netherlands) and Hungary. *Scripta Geologica* 73: 1–173.
- Rey JM (2007) *Crocidura suaveolens* (Pallas, 1811). In: Palomo LJ, Gisbert J, Blanco JC (eds) *Atlas y Libro Rojo de los Mamíferos Terrestres de España, 125–127*. Dirección General para la Biodiversidad-SECEM-SECEMU, Madrid, Spain.
- Rey JM, Landin A (1973) Sobre la presencia de *Crocidura suaveolens* en el sur de Andalucía (Mammalia, Insectivora). *Boletín de la Real Sociedad española Historia Natural (Biología)* 71: 9–16.
- Rofes J (2023) Los Micromamíferos de la Cueva de Ardales. In: Ramos Muñoz J, Weniger GC (eds) *Las sociedades prehistóricas (del Paleolítico medio al Neolítico final) en la Cueva de Ardales y Sima de las Palomas de Teba (Málaga, España)*, 121–129. Universidad de Cádiz, Editorial UCA Cádiz, Spain.
- Rofes J, Cuenca-Bescós G (2011) Evolutionary history and biogeography of the genus *Crocidura* (Mammalia, Soricidae) in Europe, with emphasis on *Crocidura kornfeldi*. *Mammalian Biology* 76: 64–78.
- Rofes J, Zuluaga MC, Murelaga X, Fernández-Eraso J, Bailon S, Iriarte MJ, Ortega LÁ, Alonso-Olazabal A (2013) Paleoenvironmental reconstruction of the early Neolithic to middle Bronze Age Peña Larga rock shelter (Álava, Spain) from the small mammal record. *Quaternary Research* 79: 158–167.
- Rofes J, Cucchi T, Hanot P, Herman J, Stephan P, Cersoy S et al. (2018) Postglacial recolonization and Holocene diversification of *Crocidura suaveolens* (Mammalia, Soricidae) on the north-western fringe of the European continent. *Quaternary Science Reviews* 190: 1–10.
- Rofes J, Ordiales A, Iriarte E, Cuenca-Bescós G, Galindo-Pellicena MÁ, Pérez-Romero A, Carretero JM, Arsuaga JL (2021) Human activities, biostratigraphy and past environment revealed by small-mammal associations at the chalcolithic levels of El Portalón de Cueva mayor (Atapuerca, Spain). *Quaternary* 4: 16.
- Román J, Ruiz G (2003) Nuevos datos sobre la distribución de la musaraña de campo *Crocidura suaveolens* (Pallas, 1811) en el sur de la península ibérica.

- Rzebik-Kowalska B (1998) Fossil history of shrews in Europe. In: Wójcik JM, Wolsan M (eds) *Evolution of Shrews*, 23–92. Polish Academy of Sciences, Białowieża, Poland.
- Saint-Girons M-C, Fons R, Nicolau-Guillaumet P (1979) Caractères distinctifs de *Crocidura russula*, *Crocidura leucodon* et *Crocidura suaveolens* en France continentale. *Mammalia* 43: 511–518.
- Sánchez Goñi MF, Landais A, Fletcher WJ, Naughton F, Desprat S, Duprat J (2008) Contrasting impacts of Dansgaard–Oeschger events over a western European latitudinal transect modulated by orbital parameters. *Quaternary Science Reviews* 27: 1136–1151.
- Schirmacher J, Kneisel J, Knitter D, Hamer W, Hinz M, Schneider RR, Weinelt M (2020) Spatial patterns of temperature, precipitation, and settlement dynamics on the Iberian Peninsula during the Chalcolithic and the Bronze Age. *Quaternary Science Reviews* 233: 106220.
- Sesé C, Sevilla P (1996) Los micromamíferos del Cuaternario peninsular español: cronoestratigrafía e implicaciones bioestratigráficas. *Revista Española de Paleontología* No Extraordinario: 278–287.
- Sesé C, Panera Gallego J, Rubio-Jara S, Pérez-González A (2011) Micromamíferos del Pleistoceno Medio y Pleistoceno Superior en el Valle del Jarama: yacimientos de Valdocarros y HAT (Madrid, España).
- Sesé C, Soto E, Santonja Gómez M, Pérez-González A, Domínguez-Rodrigo M (2016) Los micromamíferos (Lagomorpha, Eulipotyphla y Rodentia) del yacimiento del Pleistoceno Medio de Cuesta de la Bajada (Teruel, España). Estudio sistemático y consideraciones paleoambientales. *Estudios Geológicos* 72: e057.
- Sommer RS, Nadachowski A (2006) Glacial refugia of mammals in Europe: evidence from fossil records. *Mammal Review* 36: 251–265.
- Storch G, Uerpman HP (1976) Die Kleinsäugerknochen vom Castro do Zambujal. In: *Die Fauna vom Castro do Zambujal, Studien über frühe Tierknochenfunde von der Iberischen Halbinsel*, 130–138. Institut für Paläoanatomie, Domestikationsforschung und Geschichte der Tiermedizin der Universität München (Studien über frühe Tierknochenfunde von der Iberischen Halbinsel), Munich, Germany.
- Suchéras-Marx B, Escarguel G, Ferreira J, Hammer Ø (2019) Statistical confidence intervals for relative abundances and abundance-based ratios: simple practical solutions for an old overlooked question. *Marine Micropaleontology* 151: 101751.
- Taberlet P, Fumagalli L, Wust-Saucy A-G, Cosson J-F (1998) Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* 7: 453–464.
- Valenzuela S, Poitevin F, Cornette R, Bournery A, Nadal J, Vigne J-D (2009) Evolving ecosystems: ecological data from an Iron Age small mammal accumulation at Alorda Park (Catalonia, Spain). *Journal of Archaeological Science* 36: 1248–1255.
- Vigne J-D (1992) Zooarchaeology and the biogeographical history of the mammals of Corsica and Sardinia since the last ice age. *Mammal Review* 22(2): 87–96.
- Walker M, Head MJ, Berkelhammer M, Björck S, Cheng H, Cwynar L et al. (2018) Formal ratification of the subdivision of the Holocene Series/Epoch (Quaternary System/Period): two new Global Boundary Stratotype Sections and Points (GSSPs) and three new stages/subseries. *Episodes* 41(4): 213–223.
- Wilson DE, Mittermeier RA (2018) *Handbook of the Mammals of the World*. Lynx Edicions, Barcelona, Spain.
- Woods R, Marr MM, Brace S, Barnes I (2017) The small and the dead: a review of ancient DNA studies Analysing micromammal species. *Genes* 8: 312.
- Zabala J (1985) Distribución y biometría del género *Crocidura* (Wagler, 1832) en el País Vasco. *Sociedad de Estudios Vascos Cuadernos de Ciencias Naturales* 21: 71–94.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's website.

Appendix S1. Measurements of the upper fourth premolar (P4) employed for the biometric characterization.